

考試科目	計算機數學	系所別	資訊科學系 資訊安全碩士學位學程	考試時間	2月3日(五)第二節
------	-------	-----	---------------------	------	------------

本次考試共 25 題單選題，每題 4 分。

選擇題請在答案卡上作答，否則不予計分。

1. If $\begin{bmatrix} 11 & 5 \\ 35 & 16 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ c & 1 \end{bmatrix}$ and $a, b, c \in R$, then $a + b + c = ?$

(A) 7 (B) 8 (C) 9 (D) 10

2. How many of the following statements are true?

- If E is an elementary matrix, then $\det(E) = \pm 1$.
- For any $A, B \in M^{n \times n}(F)$, $\det(AB) = \det(A) \cdot \det(B)$.
- A matrix $A \in M^{n \times n}(F)$ has rank n if and only if $\det(A) \neq 0$.
- For any $A \in M^{n \times n}(F)$, $\det(A^t) = -\det(A)$.

(A) 0 (B) 1 (C) 2 (D) 3 (E) 4

3. Let A be an $m \times n$ matrix whose null space has dimension k . Which conclusion is correct?

- (A) The dimension of $NULL(A^T)$ is k .
- (B) The dimension of row space of A is $m - k$.
- (C) The dimension of column space of A is $m - k$.
- (D) The dimension of row space of A is $n - k$.

4. How many of the following vector functions are linear transformations?

● $T_1\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x^2 \\ x + y \\ y^2 \end{bmatrix}$

● $T_2\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} x + y \\ x + y + z \\ 0 \end{bmatrix}$

● $T_3\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} e^{x+y} \\ \sqrt{y} \end{bmatrix}$

● $T_4\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x+y \\ 10 \end{bmatrix}$

(A) 0 (B) 1 (C) 2 (D) 3 (E) 4

考試科目	計算機數學	系所別	資訊科學系 資訊安全碩士學位學程	考試時間	2月3日(五)第二節
------	-------	-----	---------------------	------	------------

5. How many of the following statements are true?

- The Gram–Schmidt orthogonalization process allows us to construct an orthonormal set from an arbitrary set of vectors.
- An orthonormal basis must be an ordered basis.
- Every orthogonal set is linearly independent.
- Every orthonormal set is linearly independent

(A)0 (B)1 (C)2 (D)3 (E)4

6. Let $A = \begin{bmatrix} 2 & -1 \\ -2 & 3 \end{bmatrix}$, please find A^{100}

(A) $\begin{bmatrix} -4^{100} & 1 - 4^{100} \\ 0 & 1 \end{bmatrix}$ (B) $\begin{bmatrix} 4^{100} & 1 - 4^{100} \\ 0 & 1 \end{bmatrix}$ (C) $\begin{bmatrix} 4^{100} & 1 - 4^{100} \\ 0 & -1 \end{bmatrix}$ (D) $\begin{bmatrix} 4^{100} & 1 + 4^{100} \\ 0 & 1 \end{bmatrix}$

7. How many of the following statements are true?

- Every linear operator on an n -dimensional vector space has n distinct eigenvalues.
- Any two eigenvectors are linearly independent.
- Similar matrices always have the same eigenvalues.
- Similar matrices always have the same eigenvectors.

(A)0 (B)1 (C)2 (D)3 (E)4

For problems 8-10, please find a singular value decomposition for the following matrix.

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix} = U\Sigma V$$

8. $U = ?$

(A) $\begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{\sqrt{2}}{\sqrt{3}} & 0 \\ \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ (B) $\begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{\sqrt{2}}{\sqrt{3}} & 0 \\ \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & \frac{-1}{\sqrt{2}} \end{bmatrix}$ (C) $\begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{\sqrt{2}}{\sqrt{3}} & 0 \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ (D) $\begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{\sqrt{2}}{\sqrt{3}} & 0 \\ \frac{-1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{2}} \\ \frac{-1}{\sqrt{3}} & \frac{-1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \end{bmatrix}$

考試科目	計算機數學	系所別	資訊科學系 資訊安全碩士學位學程	考試時間	2月3日(五)第二節
------	-------	-----	---------------------	------	------------

9. $\Sigma = ?$

(A) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \sqrt{3} & 0 \\ 0 & 0 & \sqrt{3} \end{bmatrix}$ (B) $\begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & \sqrt{3} & 0 \\ 0 & 0 & \sqrt{3} \end{bmatrix}$ (C) $\begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{3} \end{bmatrix}$ (D) $\begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & \sqrt{3} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

10. $V = ?$

(A) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{3}{\sqrt{2}} & \frac{3}{\sqrt{2}} \\ 0 & \frac{3}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{bmatrix}$ (B) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{3}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{bmatrix}$ (C) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{bmatrix}$ (D) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{-3}{\sqrt{2}} \end{bmatrix}$

11. Determine whether each of these compound propositions is satisfiable.

- (1) $(p \vee \neg q) \wedge (\neg p \vee q) \wedge (\neg p \vee \neg q)$.
 (2) $(p \rightarrow q) \wedge (p \rightarrow \neg q) \wedge (\neg p \rightarrow q) \wedge (\neg p \rightarrow \neg q)$.
 (3) $(p \vee q) \wedge (\neg p \vee q) \wedge (\neg p \vee \neg q)$.
 (A)(1) (B)(1), (2) (C)(2), (3) (D)(1), (3)

12. Let $S = \{a, \{a\}, \phi, \{\phi\}\}$, and $P(S)$ denote the power set of S . How many of the following statements are true?

- $a \in S$
- $\{a\} \subseteq S$
- $\{\{a\}\} \subseteq S$
- $\phi \in S$
- $\phi \subseteq S$
- $\phi \in P(S)$
- $\phi \subseteq P(S)$
- $\{\phi\} \in P(S)$
- $\{\phi\} \subseteq P(S)$


(A)6 (B)7 (C)8 (D)9

考試科目	計算機數學	系所別	資訊科學系 資訊安全碩士學位學程	考試時間	2月3日(五)第二節
------	-------	-----	---------------------	------	------------

13. Let set $A = \{1,2,3,4\}$. Define a relation of R of A as $R = \{(1,3), (1,4), (2,3), (2,4), (3,1), (3,4)\}$. Which of the following properties does this relation have?

- (1) symmetric
 - (2) asymmetric
 - (3) antisymmetric
 - (4) reflexive
 - (5) irreflexive
 - (6) transitive
- (A)(1), (4), (6) (B)(2), (5), (6) (C)(3), (5) (D)(5)

14. Consider a graph



How many of the following statements are true?

- It is bipartite.
- It has the longest simple path of length 8.
- It has an Euler circuit.
- It doesn't have an Euler circuit.

(A)0 (B)1 (C)2 (D)3

15. How many of the following statements are true?

- A graph G has a spanning tree if G is connected.
- A graph $G = (V, E)$ with $|E| = m$ satisfying $2m = \sum_{v \in V} \deg(v)$.
- A graph $G = (V_1, V_2, E)$ is bipartite, when G has a Hamilton cycle, $|V_1| = |V_2|$.
- A graph $G = (V_1, V_2, E)$ is bipartite, when G has a Hamilton cycle, $||V_1| - |V_2|| \leq 1$.

(A)1 (B)2 (C)3 (D)4

For problems 16-18, please solve the linear recurrence relation $a_n + 6a_{n-1} + 9a_{n-2} = (-3)^n$ with $a_0 = 2$ and $a_1 = 3$, and let $a_n = (i + jn + kn^2) \cdot (-3)^n$.

16. $i = ?$ (A)1 (B)2 (C)-2 (D)3

考試科目	計算機數學	系所別	資訊科學系 資訊安全碩士學位學程	考試時間	2月3日(五)第二節
------	-------	-----	---------------------	------	------------

17. $j = ?$ (A) $-\frac{7}{2}$ (B) $\frac{7}{2}$ (C) $-\frac{5}{2}$ (D) $\frac{5}{2}$

18. $k = ?$ (A) $\frac{3}{2}$ (B) $-\frac{3}{2}$ (C) $-\frac{1}{2}$ (D) $\frac{1}{2}$

For problems 19-20, please find $X = 101^{-1}$ modulo 4620.

Let $X = 100 \cdot a + b$.

19. $a = ?$ (A)15 (B)16 (C)17 (D)18

20. $b = ?$ (A)0 (B)1 (C)2 (D)3

For problems 21-22, suppose E and F are events in a sample space with $p(E) = \frac{1}{3}$, $p(F) = \frac{1}{2}$,

and $p(E|F) = \frac{2}{5}$. Find $p(F|E) = \frac{a}{b}$.

21. $a = ?$ (A)1 (B)2 (C)3 (D)4

22. $b = ?$ (A)2 (B)3 (C)4 (D)5

23. Which the following statement is false?

- (A) If NFA with k states accepts any character at all, then it cannot accept a string of length $< k$
- (B) The set for all the string that does not belong to a particular regular language L , is also a regular language
- (C) The result of subset operation of a regular language set can still be regular
- (D) Any kind of NFA can always convert to a DFA

24. Let N be an NFA with n states, let k be the number of states of a minimal DFA which is equivalent to N . Which one of the following is necessarily true?

- (A) $k \geq n^2$
- (B) $k \geq 2^n$
- (C) $k \leq n^2$
- (D) $k \leq 2^n$

25. Which of the following is not context-free language?

- (A) $L1 : \{ 0^p 1^q 0^r \mid p = q \text{ and } pqr \geq 0 \}$
- (B) $L2 : \{ 0^p 1^q 0^r \mid p = q = r \text{ and } pqr \geq 0 \}$
- (C) $L1 : \{ 0^p 1^q 0^r \mid p = q \text{ or } q = r \text{ and } pqr \geq 0 \}$
- (D) all of above are context-free language

備

註

- 一、作答於試題上者，不予計分。
- 二、試題請隨卷繳交。