



# A MATHEMATICAL MODEL OF ENTERPRISE COMPETITIVE ABILITY AND PERFORMANCE THROUGH A PARTICULAR EMDEN-FOWLER EQUATION\*

Pai Jente<sup>1</sup> Li Mengrong<sup>2†</sup> Chang Yueloong<sup>3</sup> Chiu Sumiao<sup>2</sup>

1. Department of Land Economics, National Chengchi University, Taipei 116, Taiwan, China

E-mail: [brianpai@nccu.edu.tw](mailto:brianpai@nccu.edu.tw)

2. Department of Mathematical Sciences, National Chengchi University, Taipei 116, Taiwan, China

E-mail: [liwei@math.nccu.edu.tw](mailto:liwei@math.nccu.edu.tw); [hierteliwei@gmail.com](mailto:hierteliwei@gmail.com)

3. Department of Psychology, National Chengchi University, Taipei 116, Taiwan, China

E-mail: [cyl.88054@gmail.com](mailto:cyl.88054@gmail.com)

**Abstract** In this paper we work with the ordinary differential equation  $u'' - u^3 = 0$  and obtain some interesting phenomena concerning blow-up, blow-up rate, life-span, zeros and critical points of solutions to this equation.

**Key words** estimate; life-span; blow-up; blow-up rate; performance; competitive ability

**2000 MR Subject Classification** 34A34; 34C05; 34C11; 34C99

## 0 Introduction

How to improve the performance and competitiveness of the company is the critical issue of Industrial and Organizational Psychology in Taiwan. We try to design an appropriate mathematical model of the competitiveness and the performance of the 293 benchmark enterprises out of 655 companies.

Unexpectedly, we discover the correlation of performance and competitiveness is extremely high. Some benchmark enterprises present the following phenomena.

Competitive ability (force,  $F(P(n))$ ) is a cubic function of the performance ( $P(n)$ ); that is, there exist positive constant performance  $P_0 > 0$ , and a positive constant  $k > 0$  such that

$$F(P(n)) = k(P(n) - P_0)^3,$$

where  $n$  is the surveying rod enterprise's composition department number or the main unit commanders counts, the performance  $P(n)$  of the rod enterprise is larger than  $P_0$  and  $F$  is proportional to the second-order derivative of  $P$  with respect to  $n$ .

\*Received June 23, 2010. This work is financed by NSC, Metta Education, Grand Hall Company and Auria Solar Company.

†Corresponding author: Li Mengrong.

Although the relation between  $F(P(n))$  and  $P(n)$  is non-linear but not the same as the previous type, there exist positive constant performances  $P_0 > 0$ ,  $P_1 > 0$  and a positive constant  $k(n) > 0$  such that

$$F(P(n)) = k(P(n) - P_1)(P(n) - P_0)^2.$$

$F$  and the  $P$  relations present the following punishment type, there exist positive constant performances  $P_0 > 0$ ,  $P_1 > 0$ ,  $P_2 > 0$  and a positive constant  $k(n) > 0$  such that

$$F(P(n)) = k(P(n) - P_2)(P(n) - P_1)(P(n) - P_0).$$

Now we consider the special case 1. For  $F(P(n)) = M \frac{d^2 P(n)}{dn^2}$ , let  $u(n) := \sqrt{\frac{k}{M}}(P(n) - P_0) \geq 0$ , then we obtain a stationary one-dimensional semilinear wave equation with initial condition

$$\begin{cases} u'' - u^3 = 0, & n \geq n_0, \\ u(n_0) = u_0 = \sqrt{\frac{k}{M}}(P(n_0) - P_0), & u'(n_0) = u_1. \end{cases} \quad (0.1)$$

It is clear that the function  $u^3$  is locally Lipschitz, hence by the standard theory, the local existence of classical solutions is applicable to problem (0.1).

We would use our methods used in [1–11] to discuss problem (0.1) in two parts: blow-up time and “the singularity and regularity of solution for higher order derivatives”.

In Section 1, we would deal with the estimates for the existence interval of the solutions of (0.1); in Section 2, with the blow-up rate and blow-up constant; in Section 3, with the global existence, critical point and the asymptotic behavior; in Section 4, with the null points (zero) and triviality; in Section 5 with the stability and instability. In Sections 6 and 7, we would study the blow-up behavior of  $u^{(k)}$  and the regularity of the solution  $u$  of (0.1).

**Notations and Fundamental Lemmas** For a given function  $u$  in this work, we use the following abbreviations:

$$a_u(n) = u(n)^2, \quad E_u(n_0) = u_1^2 - \frac{1}{2}u_0^4, \quad J_u(n) = a_u(n)^{-\frac{1}{2}}.$$

**Definition** A function  $g : \mathbb{R} \rightarrow \mathbb{R}$  with a blow-up rate  $q$  means that  $g$  exists only in a finite time, that is, there is a finite number  $T^*$  such that

$$\lim_{t \rightarrow T^*} g(t)^{-1} = 0 \quad (0.2)$$

and there exists a non-zero  $\beta \in \mathbb{R}$  with

$$\lim_{t \rightarrow T^*} (T^* - t)^q g(t) = \beta, \quad (0.3)$$

in this case  $\beta$  is called the blow-up constant of  $g$ .

According to the uniqueness of the solutions to (0.1), we can rewrite  $a_u(n) = a(n)$ ,  $J_u(n) = J(n)$  and  $E_u(n) = E(n)$ . After some elementary calculations we obtain the following:

**Lemma 1** Suppose that  $u$  is the solution of (0.1), then we have

$$E(n) = u'(n)^2 - \frac{1}{2}u(n)^4 = E(n_0), \quad (0.4)$$

$$6u'(n)^2 = 4E(n_0) + a''(n), \tag{0.5}$$

$$J''(n) = 2E(n_0)J(n)^3, \tag{0.6}$$

$$J'(n)^2 = J'(n_0)^2 - E(n_0)J(n_0)^4 + E(n_0)J(n)^4, \tag{0.7}$$

and

$$a'(n) = a'(n_0) + 2E(n_0)(n - n_0) + 3 \int_{n_0}^n u(r)^4 dr. \tag{0.8}$$

The following lemmas are easy to prove, so we omit their proofs.

**Lemma 2** Suppose that  $r$  and  $s$  are real constants and  $u \in C^2(\mathbb{R})$  satisfies

$$\begin{aligned} u'' + ru' + su &\leq 0, \quad u \geq 0, \\ u(0) = 0, \quad u'(0) &= 0, \end{aligned}$$

then  $u$  must be null:  $u \equiv 0$ .

**Lemma 3** If  $g(t)$  and  $h(t, r)$  are continuous with respect to their variables and

$$\lim_{t \rightarrow T} \int_0^{g(t)} h(t, r) dr$$

exists, then

$$\lim_{t \rightarrow T} \int_0^{g(t)} h(t, r) dr = \int_0^{g(T)} h(T, r) dr.$$

## 1 Estimates for the Life-Spann

To estimate the existence interval of the solution of (0.1), we separate this section into three parts:  $E(n_0) < 0$ ,  $E(n_0) = 0$  and  $E(n_0) > 0$ . Here the existence interval  $N$  of  $u$  means that  $u$  exists and makes sense only in the interval  $[n_0, N)$  such that problem (0.1) possesses the solution  $u \in \bar{C}^2(n_0, N)$ .

### 1.1 Estimates for the Existence Intervals Under $E(n_0) \leq 0$

We deal with two cases,  $E(n_0) < 0$ , and  $E(n_0) = 0$  and  $a'(n_0) > 0$  in this subsection, but the case  $E(n_0) = 0$  and  $a'(n_0) \leq 0$  will be considered in Sections 3 and 4. Here we have the following result.

**Theorem 4** If  $N$  is the existence interval of the solution  $u$  to (0.1) with  $E(n_0) < 0$ , then  $N$  is finite. Further, for  $a'(n_0) \geq 0$  we have the estimate

$$N \leq N_1^* = n_0 + \int_0^{J(n_0)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} \tag{1.1}$$

for  $a'(n_0) < 0$ ,

$$N \leq N_2^* = n_0 + \left( \int_0^{\left(\frac{1}{2} \frac{-1}{E(n_0)}\right)^{\frac{1}{4}}} + \int_{J(n_0)}^{\left(\frac{1}{2} \frac{-1}{E(n_0)}\right)^{\frac{1}{4}}} \right) \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}}, \tag{1.2}$$

where  $\alpha = \left(\frac{1}{2} \frac{-1}{E(n_0)}\right)^{\frac{1}{4}}$ . Furthermore, if  $E(n_0) = 0$  and  $a'(n_0) > 0$ , then

$$N \leq N_3^* := n_0 + \frac{2a(n_0)}{a'(n_0)}. \quad (1.3)$$

**Proof** For  $E(n_0) < 0$ , we know that  $a(n_0) > 0$ ; otherwise we would get  $a(n_0) = 0$ , that is,  $u_0 = 0$ , then  $E(n_0) = u_1^2 \geq 0$ , this contradicts  $E(n_0) < 0$ . In this situation we separate the proof of this Theorem into two subcases,  $a'(n_0) \geq 0$  and  $a'(n_0) < 0$ .

(i)  $a'(n_0) \geq 0$ . By (0.5) and (0.7), we find that

$$\begin{cases} a'(n) \geq a'(n_0) - 4E(n_0)(n - n_0) & \forall n \geq n_0, \\ a(n) \geq a(n_0) + a'(n_0)n - 2E(n_0)(n - n_0)^2 & \forall n \geq n_0, \end{cases} \quad (1.4)$$

$$J'(n) = -\sqrt{\frac{1}{2} + E(n_0)J(n)^4} \leq J'(n_0) \quad \forall n \geq n_0 \quad (1.5)$$

and  $J(n) \leq a(n_0)^{-\frac{1}{2}} - \frac{1}{2}a(n_0)^{-\frac{3}{2}}a'(n_0)(n - n_0) \quad \forall n \geq n_0$ . Thus, there exists a finite number  $N_1^*(u_0, u_1) \leq n_0 + \frac{2a(n_0)}{a'(n_0)} = n_0 + \frac{u_0}{u_1}$  such that  $J(N_1^*(u_0, u_1)) = 0$  and so  $a(n) \rightarrow \infty$  as  $n \rightarrow N_1^*(u_0, u_1)$ . This means that  $N \leq N_1^*(u_0, u_1)$ . Now we estimate  $N_1^*(u_0, u_1)$ . By (1.5) and  $J(N_1^*(u_0, u_1)) = 0$  we find that

$$\int_{J(n)}^{J(n_0)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = n - n_0, \quad \forall n \geq n_0 \quad (1.6)$$

and hence we get estimate (1.1).

(ii)  $a'(n_0) < 0$ . By (1.4),  $a'(n_0) < 0$  and the convexity of  $a$ , we can find a unique finite number  $n_1 = n_1(u_0, u_1)$  such that

$$\begin{cases} a'(n) < 0 = a'(n_1) & \text{for } n \in (n_0, n_1), \\ a'(n) > 0 & \text{for } n > n_1, \end{cases} \quad (1.7)$$

and  $a(n_1) > 0$ . If not, then  $u(n_1) = 0$ , thus  $E(n) = E(n_1) = u'(n_1)^2 \geq 0$ ; yet this is a contradiction to  $E(n_0) < 0$ . Hence, we conclude that  $a(n) > 0$ ,  $\forall n \geq n_0$ ,  $u'(n_1) = 0$ ,  $E(n_0) = -\frac{1}{2}u(n_1)^4$  and  $J(n_1)^4 = \frac{-1}{2E(n_1)}$ .

After arguments similar to the step (i), there exists an  $N_2^* := N_2^*(u_0, u_1)$  such that the life-span  $N$  of  $u$  is bounded by  $N_2^*$ , that is,  $N \leq N_2^*$ . By an analogous argument, using (1.7), (0.7) and the fact that  $J(n_1)^4 = \frac{-1}{2E(n_0)}$  and  $J(N_2^*) = 0$ , we conclude that

$$J'(n)^2 = -E(n_0) \left( J(n_1)^4 - J(n)^4 \right), \quad \forall n \geq n_1,$$

$$J'(n)^2 = E(n_0) \left( J(n_1)^4 - J(n)^4 \right), \quad \forall n \in [n_0, n_1],$$

$$J'(n) = -\sqrt{\frac{1}{2} + E(n_0)J(n)^4}, \quad \forall n \geq n_1, \quad (1.8)$$

$$J'(n) = \sqrt{\frac{1}{2} + E(n_0)J(n)^4}, \quad \forall n \in [n_0, n_1], \quad (1.9)$$

$$\int_{J(n)}^{J(n_1)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = n - n_1, \quad \forall n \geq n_1, \tag{1.10}$$

$$\int_{J(n_0)}^{J(n_1)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = n_1 - n_0, \tag{1.11}$$

and

$$N_2^* = n_1 + \int_0^{\left(\frac{-1}{2E(n_0)}\right)^{\frac{1}{4}}} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}}. \tag{1.12}$$

Estimate (1.12) is equivalent to (1.2).

(iii) For  $E(n_0) = 0$ , by (0.6) and  $a'(n_0) > 0$ , we get that  $J'(n_0) < 0, J''(n) = 0$  and  $J(n) = a(n_0)^{-\frac{3}{2}} \left( a(n_0) - \frac{1}{2}a'(n_0)(n - n_0) \right), \forall n \geq n_0$ . Thus we conclude that

$$a(n) = a(n_0)^3 \left( a(n_0) - \frac{1}{2}a'(n_0)(n - n_0) \right)^{-2}, \quad \forall n \geq n_0 \tag{1.13}$$

and (1.3) is proved.

### 1.2 Estimates for the Life-Spann under $E(n_0) > 0$

In this subsection we consider the case  $E(n_0) > 0$ . We have the following blow-up result.

**Theorem 5** If  $N^*$  is the existence interval of  $u$  which solves problem (0.1) with  $E(n_0) > 0$ , then  $N^*$  is finite. Further, in case of  $a'(n_0) > 0$ , we have

$$N^* \leq N_4^*(u_0, u_1) = n_0 + \int_0^{J(n_0)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}}. \tag{1.14}$$

In the case of  $a'(n_0) = 0$  we have

$$N^* \leq N_5^*(u_0, u_1) = n_0 + \int_0^\infty \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}}. \tag{1.15}$$

For  $a'(n_0) < 0$  and  $z(u_0, u_1)$  given by

$$z(u_0, u_1) = n_0 + \int_0^{\sqrt{a(n_0)}} \frac{dr}{\sqrt{E(n_0) + \frac{1}{2}r^4}} \tag{1.16}$$

is the zero of  $a$ . Further we have

$$N^* \leq N_6^*(u_0, u_1) := (z + N_5^*)(u_0, u_1). \tag{1.17}$$

**Proof** The case of a zero for  $u$  is postponed to Section 4.

i) For  $a'(0) > 0$ , by (0.6) we have

$$\begin{cases} mJ''(n) = (mJ(n))^3, \\ mJ(n_0) = ma(n_0)^{-\frac{1}{2}}, \quad mJ'(n_0) = \frac{-1}{2}ma(n_0)^{-\frac{3}{2}}a'(n_0), \end{cases}$$

where  $m := (2E(n_0))^{\frac{1}{2}}$ . Now we set

$$\tilde{E}(n) := 2E(n_0)J'(n)^2 - 2E(n_0)^2(mJ(n))^4. \tag{1.18}$$

From some calculations, we see that  $\tilde{E}(n)$  is a constant and by using (0.8) we obtain that

$$\tilde{E}(n) = \frac{1}{2}m^2 = \tilde{E}(n_0) = E(n_0), \quad (1.19)$$

$$\frac{1}{2} = J'(n)^2 - E(n_0)J(n)^4,$$

$$a'(n) \geq a'(n_0) + 2E(n_0)(n - n_0) > 0 \quad \forall n \geq n_0, \quad (1.20)$$

$$J'(n) < 0 \quad \forall n \geq n_0,$$

$$J'(n) = -\sqrt{\frac{1}{2} + E(n_0)J(n)^4} \quad \forall n \geq n_0, \quad (1.21)$$

and

$$\int_{J(n)}^{J(n_0)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = n - n_0 \quad \forall n \geq n_0. \quad (1.22)$$

By (1.21), there exists a finite number  $N_4^*(u_0, u_1)$  such that  $J(N_4^*(u_0, u_1)) = 0$ , and from (1.22), estimate (1.14) follows easily.

ii) From  $a'(n_0) = 0 = u_0$ ,  $E(n_0) = u_1^2$  and (0.8) we obtain

$$a'(n) = 2E(n_0)n + 3 \int_{n_0}^n u(r)^4 dr \quad \forall n \geq n_0, \quad (1.23)$$

$$a(n) > 0 \quad \forall n \geq n_0,$$

thus  $J(n)$  can be defined for each  $n > n_0$  and  $J'(n) < 0$ ,  $\forall n > n_0$ .

Using (0.6), we conclude that, for each  $\tilde{n} > n_0$ ,

$$J'(n) = -\sqrt{J'(\tilde{n})^2 - E(n_0)(J(\tilde{n})^4 - J(n)^4)}, \quad \forall n \geq \tilde{n}, \quad (1.24)$$

$$\lim_{\tilde{n} \rightarrow 0} J'(\tilde{n})^2 - u_1^2 J(\tilde{n})^4 = \frac{1}{2}, \quad (1.25)$$

thus after inducing (1.24) and (1.25), estimate (1.15) follows.

(iii) For  $a'(n_0) < 0$ , by (1.20) we have  $a'(n) \geq 0$  for large  $n$ .

Suppose  $z$  is the first positive number  $n$  so that  $a'(n) = 0$ , then  $u(z) = 0$ ; otherwise,  $u'(z) = 0$  and  $E(z) = -\frac{1}{2}u(z)^4 < 0$ , this contradicts the assumption  $E(n_0) = E(z) > 0$ . After the time  $n = z$ , same as the procedures given in the proof of (i), using (1.22) we obtain (1.17).

### 1.3 Some Properties Concerning the Life-Spann $N_1^*(u_0, u_1)$


In principle,  $N_1^*(u_0, u_1)$  depends on three variables  $u_0$  and  $u_1$ . Set  $c_k := \frac{2u_1^2}{u_0^4}$ , then

$$N_1^*(u_0, u_1) = \sqrt{2}u_0^{-1} (\sqrt[3]{1 - c_k})^{-1} \int_0^{\sqrt[4]{1 - c_k}} \frac{dr}{\sqrt{1 - r^4}}.$$

For convenience, we consider the case  $u_1 = 0$ ,

$$N_1^*(u_0, 0) = \frac{\sqrt{\pi}}{2\sqrt{2}} u_0^{-1} \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})}.$$

Using Maple we obtain the graphs of  $N_1^*(u_0, 0)$  below:



To grasp the property of the existence interval  $N_1^* := N_1^*(u_0, u_1)$  is very difficult, after some computations we get

$$N_1^* = \sqrt{2} (u_0^4 - 2u_1^2)^{-\frac{1}{4}} \int_0^{\sqrt[4]{1 - \frac{2}{u_0^4} u_1^2}} \frac{dr}{\sqrt{1 - r^4}}.$$

By the experience of studying the existence interval  $N_1^*$ , we consider its properties with  $a'(n_0) \geq 0$  in three cases:

**Case 1**  $0 < u_0^4 - 2u_1^2 < 1$ . In this situation we find that

- (i) for fixed  $u_1, u_0 \frac{\partial}{\partial u_0} N_1^* < 0$ .
- (ii) for fixed  $u_0$ , the life-spann  $N_1^*(u_0, u_1)$  decreases in  $u_1^2$ .

**Case 2**  $u_0^4 - 2u_1^2 > 1$ . The life-spann  $N_1^*(u_0, u_1) = N_1^*(s, u_0)$  decreases in  $s = \sqrt[4]{1 - \frac{2}{u_0^4} u_1^2}$  for fixed  $u_0$ .

**Case 3**  $u_0^{p+1} - 2u_1^2 = 1$ . On the region  $\{(u_0, u_1) \in \mathbb{R}^2 \mid u_0^4 - 2u_1^2 = 1\}$ , we find that

$$N_1^*(u_0, u_1) = N_1^*(u_0) = \sqrt{2} \int_0^{u_0^{-1}} \frac{1}{\sqrt{1 - r^4}} dr,$$

and that  $N_1^*(u_0)$  is monotone decreasing in  $u_0$ .

## 2 Blow-up Rate and Blow-up Constant

In this section we study the blow-up rate and blow-up constant for  $a, a'$  and  $a''$  under the conditions in Section 1. We have got the following results.

**Theorem 6** If  $u$  is the solution of the problem (0.1) with one of the following properties:

- (i)  $E(n_0) < 0$ ,
- (ii)  $E(n_0) = 0, a'(n_0) > 0$ ,
- (iii)  $E(n_0) > 0$ .

Then the blow-up rate of  $a$  is 2, and the blow-up constant  $K_1$  of  $a$  is 2, that is, for  $m = 1, 2, 3, 4, 5, 6$ ,

$$\lim_{n \rightarrow N_m^*} (N_m^* - n)^2 a(n) = 2. \tag{2.1}$$

The blow-up rate of  $a'$  is 3, and the blow-up constant  $K_2$  of  $a'$  is 4, that is, for  $m = 1, 2, 3, 4, 5, 6$ ,

$$\lim_{n \rightarrow N_m^*} (N_m^* - n)^3 a'(n) = 4. \tag{2.2}$$

The blow-up rate of  $a''$  is 4, and the blow-up constant  $K_3$  of  $a''$  is 12, that is, for  $m = 1, 2, 3, 4, 5, 6$ ,

$$\lim_{n \rightarrow N_m^*} a''(n) (N_m^* - n)^4 = 12. \quad (2.3)$$

**Proof** (i) Under this condition,  $E(n_0) < 0$ ,  $a'(n_0) \geq 0$  by (1.1), (1.6) and Lemma 4, we get

$$\int_0^{J(n)} \frac{1}{N_1^* - n} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = 1 \quad \forall n \geq n_0, \quad (2.4)$$

$$\lim_{n \rightarrow N_1^*} \sqrt{2} \frac{J(n)}{N_1^* - n} = 1. \quad (2.5)$$

Identity (2.5) is equivalent to (2.1) for  $m = 1$ .

For  $E(n_0) < 0$ ,  $a'(n_0) < 0$ , by (1.9) we have also

$$\int_0^{J(n)} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = N_2^* - n, \quad \forall n \geq n_0. \quad (2.6)$$

Through Lemma 4 and (2.6), therefore we get (2.1) for  $m = 2$ .

Seeing (1.5) and (1.8), we find

$$\lim_{n \rightarrow N_m^*} J'(n) = -\frac{1}{2}\sqrt{2}, \quad (2.7)$$

$$\lim_{n \rightarrow N_m^*} a'(n) (N_m^* - n)^3 = 4, \quad (2.8)$$

$$\lim_{n \rightarrow N_m^*} u'(n)^2 (N_m^* - n)^4 = 2 \quad (2.9)$$

for  $m = 1, 2$ . Using (0.5) and (2.9) we obtain, for  $m = 1, 2$ ,

$$\lim_{n \rightarrow N_m^*} a''(n) (N_m^* - n)^4 = 6 \lim_{n \rightarrow N_m^*} u'(n)^2 (N_m^* - n)^4. \quad (2.10)$$

Thus, (2.10) and (2.3) are equivalent.

(ii) For  $E(n_0) = 0$ ,  $a'(n_0) > 0$ , by (1.1) we get, for  $m = 1, 2$ ,

$$a(n) = a(n_0)^3 \left( \frac{1}{2} a'(n_0) \right)^{-2} \cdot (N_3^* - n) \quad \forall n \geq n_0. \quad (2.11)$$

Therefore, estimates (2.1), (2.2) and (2.3) for  $m = 3$  are followed from (2.11).

(iii) For  $E(n_0) > 0$ , estimates (2.1), (2.2) and (2.3) for  $m = 4, 5, 6$ , are similar to the above arguments (i) in the proof of this theorem.

### 3 Global Existence and Critical Point

In this section we study the following case that  $E(n_0) = 0$  and  $a'(n_0) < 0$ .

Here we take the global existence of the solutions to the problem (0.1) in the following sense:

$$J(n) > 0, \quad a'(n)^{-2} > 0, \quad a''(n)^{-2} > 0 \quad \forall n \in [n_0, N],$$

where  $N$  is the time that  $u$  exists, in other words, in any finite time  $u$  does not blow up in  $C^2$  sense, even though  $u$  blows up in a finite time in some sense, for example,  $C^k$  or  $L^k$  for some  $k \geq 3$ .

By [12, p.151] every positive proper solution of problem (0.1) has the asymptotic form

$$u(n) \sim cn^{-1}.$$

This result could happen and will be explained below only in the case that  $E(n_0) = 0$  and  $a'(n_0) < 0$ . Under the condition it is easy to see that  $J(n) > 0 \forall n \in (n_0, N)$ , and

$$a(n) = a(n_0)^3 \left( a(n_0) - \frac{1}{2}a'(n_0)(n - n_0) \right)^{-2}, \quad \forall n \in (n_0, N),$$

$$a'(n)^{-2} = a(n_0)^{-6} a'(n_0)^{-2} \left( a(n_0) - \frac{1}{2}a'(n_0)(n - n_0) \right)^6 > 0, \quad \forall n \in (n_0, N),$$

$$a''(n)^{-2} = \frac{4}{9}a(n_0)^{-6} a'(n_0)^{-4} \left( a(n_0) - \frac{1}{2}a'(n_0)(n - n_0) \right)^8 > 0, \quad \forall n \in (n_0, N).$$

Hence we find  $\lim_{n \rightarrow \infty} a(n) = 0, \lim_{n \rightarrow \infty} a'(n) = 0, \lim_{n \rightarrow \infty} a''(n) = 0$  and

$$\lim_{n \rightarrow \infty} (n - n_0)^2 a(n) = a(n_0)^3 \left( \frac{1}{-2}a'(n_0) \right)^{-2} = \frac{u_0^4}{u_1^2}, \tag{3.1}$$

$$\lim_{n \rightarrow \infty} (n - n_0)^3 a'(n) = a(n_0)^3 a'(n_0) \left( \frac{1}{-2}a'(n_0) \right)^{-3} = -\frac{2u_0^4}{u_1^2}, \tag{3.2}$$

$$\lim_{n \rightarrow \infty} (n - n_0)^4 a''(n) = \frac{3}{2}a(n_0)^3 a'(n_0)^2 \left( \frac{1}{-2}a'(n_0) \right)^{-4} = \frac{3}{32} \frac{u_0^4}{u_1^2}. \tag{3.3}$$

**Theorem 7** Suppose that  $u$  is the solution of problem (0.1) with  $E(n_0) = 0$  and  $a'(n_0) < 0$ , then  $u$  can be defined globally and estimates (3.1), (3.2) and (3.3) are valid.

### 4 Existence of Zero and Triviality

In this section we discuss the triviality of the solution for problem (0.1) under the case that  $E(n_0) = 0, a'(n_0) = 0$ .

**Proposition** If  $u$  is the solution of problem (0.1) with  $E(n_0) = 0$  and  $a'(n_0) = 0$ , then  $u$  must be null.

**Proof** Under the conditions  $E(n_0) = 0, a'(n_0) = 0$ , by using (0.5), it is easy to see that  $u_0 = 0 = u_1$ , herein the supremum below exists:

$$n_1 := \sup \{ \alpha : a(n) \leq 1, \forall n \in [n_0, \alpha] \},$$

and then  $2u'(n)^2 = u(n)^4 \geq 0, a''(n) = 6u'(n)^2 = 3u(n)^4 = \frac{3}{2}a(n)^2$ . By Lemma 2 we conclude that

$$a''(n) \leq 6a(n), \quad a(n) \equiv 0 \equiv u(n) \quad \text{in } [n_0, n_1].$$

Continue these steps we get the assertion of this theorem.

For the case that  $E(n_0) > 0 > a'(n_0)$ , we have the following result.

**Theorem 8** Suppose that  $u$  is the solution to problem (0.1) with  $E(n_0) > 0 > a'(n_0)$  and  $z(u_0, u_1)$  given by

$$z(u_0, u_1) = n_0 + \int_0^{\sqrt{a(n_0)}} \frac{dr}{\sqrt{E(n_0) + \frac{1}{2}r^4}}, \tag{4.1}$$

then  $z(u_0, u_1)$  is the zero of  $a$ . Furthermore, we have

$$\lim_{n \rightarrow z^-(u_0, u_1)} a(n) \left( z(u_0, u_1) - n \right)^{-2} = E(n_0)^2, \tag{4.2}$$

$$\lim_{n \rightarrow z^-(u_0, u_1)} \left( z(u_0, u_1) - n \right)^{-1} a'(n) = -2E(n_0)^{\frac{3}{2}}, \tag{4.3}$$

$$\lim_{n \rightarrow z^-(u_0, u_1)} a''(n) = 2E(n_0). \tag{4.4}$$

**Proof** 1) For  $E(n_0) > 0 > a'(n_0)$ , by (0.4) we obtain that

$$a'(n) = -2\sqrt{E(n_0)a(n) + \frac{1}{2}a(n)^3}, \tag{4.5}$$

$$z(u_0, u_1) = n_0 + \int_0^{a(n_0)} \frac{dr}{2\sqrt{E(n_0)r + \frac{1}{2}r^3}}, \tag{4.6}$$

$$n = n_0 + \int_{a(n)}^{a(n_0)} \frac{dr}{2\sqrt{E(n_0)r + \frac{1}{2}r^3}}, \tag{4.7}$$

and

$$\begin{aligned} z(u_0, u_1) &= n_0 + \int_0^{a(n_0)} \frac{dr}{2\sqrt{r}\sqrt{E(n_0) + \frac{1}{2}r^2}} = n_0 + \int_0^{\sqrt{a(n_0)}} \frac{dr}{\sqrt{E(n_0) + \frac{1}{2}r^4}} \\ &= n_0 + 2^{\frac{1}{4}}E(n_0)^{-\frac{1}{2}} \int_0^{(2E(n_0))^{-\frac{1}{4}}\sqrt{a(n_0)}} \frac{dr}{\sqrt{1+r^4}}. \end{aligned} \tag{4.8}$$

Thus, (4.1) is proved.

2) To claim (4.2), by (4.6), (4.7) and Lemma 3, it induces that

$$\begin{aligned} z(u_0, u_1) - n &= \int_0^{a(n)} \frac{dr}{2\sqrt{E(n_0)r + \frac{1}{2}r^3}} = 2^{\frac{1}{4}}E(n_0)^{-\frac{1}{4}} \int_0^{(2E(n_0))^{-\frac{1}{4}}\sqrt{a(n)}} \frac{dr}{\sqrt{1+r^4}}, \\ (z(u_0, u_1) - n)^{-1} &\int_0^{(2E(n_0))^{-\frac{1}{4}}\sqrt{a(n)}} \frac{dr}{\sqrt{1+r^4}} = \sqrt[4]{\frac{E(n_0)}{2}}, \\ \sqrt[4]{\frac{E(n_0)}{2}} &= \lim_{n \rightarrow z^-(u_0, u_1)} (z(u_0, u_1) - n)^{-1} \int_0^{(2E(n_0))^{-\frac{1}{4}}\sqrt{a(n)}} \frac{dr}{\sqrt{1+r^4}} \\ &= \lim_{n \rightarrow z^-(u_0, u_1)} (z(u_0, u_1) - n)^{-1} (2E(n_0))^{-\frac{1}{4}} \sqrt{a(n)} \\ &\cdot \lim_{n \rightarrow z^-(u_0, u_1)} \int_0^1 \frac{ds}{\sqrt{1 + \left( (2E(n_0))^{-\frac{1}{4}} \sqrt{a(n)}s \right)^4}} \\ &= \lim_{n \rightarrow z^-(u_0, u_1)} (z(u_0, u_1) - n)^{-1} (2E(n_0))^{-\frac{1}{4}} \sqrt{a(n)}. \end{aligned} \tag{4.9}$$

Thus we get conclusion (4.2).

3) Using (4.8) and (4.5) we obtain that

$$\begin{aligned} & \lim_{n \rightarrow z^-(u_0, u_1, p)} (z(u_0, u_1, p) - n)^{-1} a'(n) \\ &= -2 \lim_{n \rightarrow z^-(u_0, u_1, p)} \sqrt{a(n) (z(u_0, u_1, p) - n)^{-2} \left( E(0) + \frac{2}{p+1} a(n)^{\frac{p+1}{2}} \right)} \\ &= -2E(0)^{\frac{3}{2}}. \end{aligned}$$

4) Applying (0.5), (4.2) and (4.3), we find

$$\begin{aligned} & \lim_{n \rightarrow z^-(u_0, u_1)} a(n) (z(u_0, u_1) - n)^{-2} a''(n) \\ &= \frac{3}{2} \lim_{n \rightarrow z^-(u_0, u_1)} (a'(n) (z(u_0, u_1) - n)^{-1})^2 - 4E(n_0) \lim_{n \rightarrow z^-(u_0, u_1)} a(n) (z(u_0, u_1) - n)^{-2} \\ &= 2E(n_0)^3. \end{aligned}$$

Hence (4.4) is proved.

### 5 Stability, Instability and Asymptotic Analysis

We now consider the applications of the theorems above to the stability theory for the problem

$$\begin{cases} u''(n) = u(n)^3, \\ u(n_0) = \varepsilon_1, u'(n_0) = \varepsilon_2. \end{cases} \tag{5.1}$$

We say problem (5.1) is stable under condition F, if any nontrivial global solution  $u \in C^2(\mathbb{R}^+)$  of (5.1) under condition F satisfies  $\|u\|_{C^2} \rightarrow 0$  for  $|\varepsilon_1| + |\varepsilon_2| \rightarrow 0$ . According to Theorems 4-8 we have the following result.

**Corollary 9.1** Problem (5.1) is stable under  $E_u(n_0) = 0$ ,  $\varepsilon_1 \varepsilon_2 < 0$ , and unstable under one of the followings:

- (i)  $E_u(n_0) < 0$ ,
- (ii)  $E_u(n_0) = 0 < \varepsilon_1 \varepsilon_2$ ,
- (iii)  $E_u(n_0) > 0$ .

According to Theorems 4 and 5, we can obtain the following conclusions, when we reconsider the problem

$$\begin{cases} u''(n) = u(n)^3, \\ u(n_0) = \varepsilon u_0, u'(n_0) = \varepsilon^2 u_1, \varepsilon > 0. \end{cases} \tag{5.2}$$

**Theorem 9.2** If  $N_\varepsilon$  is the existence interval of the solution  $u_\varepsilon$  to (5.2) with  $E(n_0) < 0$ , then  $N_\varepsilon$  is finite. Further, for  $a'(n_0) \geq 0$ , we have the estimate  $N_\varepsilon \leq N_{1,\varepsilon}^* = n_0 + \varepsilon^{-1} (N_1^* - n_0)$  for  $a'(n_0) < 0$ ,

$$N_\varepsilon \leq N_{2,\varepsilon}^* = n_0 + \varepsilon^{-1} (N_2^* - n_0).$$

Furthermore, if  $E(n_0) = 0$  and  $a'(n_0) > 0$ , then

$$N_\varepsilon \leq N_3^* := n_0 + \varepsilon^{-1} (N_2^* - n_0). \tag{5.3}$$

**Theorem 9.3** If  $N_\varepsilon^*$  is the existence interval of  $u$  which solves problem (5.2) with  $E(n_0) > 0$ , then  $N_\varepsilon^*$  is finite. Further, in case of  $a'(n_0) > 0$  we have

$$N_\varepsilon^* \leq N_{4,\varepsilon}^*(u_0, u_1) = n_0 + \varepsilon^{-1} (N_4^* - n_0).$$

In the case of  $a'(n_0) = 0$  we have

$$N_\varepsilon^* \leq N_{5,\varepsilon}^*(u_0, u_1) = n_0 + \varepsilon^{-1} (N_5^* - n_0).$$

For  $a'(n_0) < 0$ ,  $z_\varepsilon(u_0, u_1)$  given by

$$z_\varepsilon(u_0, u_1) = n_0 + \varepsilon^{-1} (z(u_0, u_1) - n_0)$$

is the zero of  $u_\varepsilon$ . Further we have

$$N_\varepsilon^* \leq N_{6,\varepsilon}^*(u_0, u_1) := (z_\varepsilon + N_{5,\varepsilon}^*)(u_0, u_1).$$

### 6 Regularity of Solution to (0.1)

In this section we study the regularity of the positive solution  $u$  of the nonlinear problem (0.1). Using (0.4) we have

$$u'(n)^2 = E(n_0) + \frac{1}{2}u(n)^4, \tag{6.1}$$

where  $E(n_0) = u_1^2 - \frac{1}{2}u_0^4$ .

#### 6.1 Regularity of Solution to (0.1)

Now consider the regularity of the positive solution  $u$  of problem (0.1). We have the following results.

**Theorem 10** If  $u$  is the positive solution of problem (0.1) with the life-spann  $N^*$ , then  $u \in C^q(n_0, N^*)$  for any  $q \in \mathbb{N}$  and

$$u^{(2k)} = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} u^{C_{k,i}}, \tag{6.2}$$

$$u^{(2k+1)} = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} C_{k,i} u^{C_{k,i}-1} u' = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} O_{k,i} u^{C_{k,i}-1} u' \tag{6.3}$$

for positive integer  $k$ , where  $\lfloor \frac{C_{k,0}}{4} \rfloor$  denotes the Gaussian integer number of  $\frac{C_{k,0}}{4}$ ,

$$C_{k,i} = 4(k-i) - 2n + 1, O_{k,i} = E_{k,i} C_{k,i}, E_{0,0} = 1,$$

$$E_{k,0} = O_{k-1,0} \left[ \frac{1}{2} (C_{k-1,0} + 1) \right] = E_{k-1,0} C_{k-1,0} \left[ \frac{1}{2} (C_{k-1,0} + 1) \right],$$

$$\begin{aligned} E_{k,k-1} &= O_{k-1,k-2} (C_{k-1,k-2} - 1) E(0) \\ &= E_{k-1,k-2} C_{k-1,k-2} (C_{k-1,k-2} - 1) E(0), \end{aligned}$$

and

$$\begin{aligned}
 E_{k,l} &= O_{k-1,l-1} (C_{k-1,l-1} - 1) E(0) + O_{k-1,l} \left[ \frac{1}{2} (C_{k-1,l} + 1) \right] \\
 &= E_{k-1,l-1} C_{k-1,l-1} (C_{k-1,l-1} - 1) E(0) + E_{k-1,l} C_{k-1,l} \left[ \frac{1}{2} (C_{k-1,l} + 1) \right]
 \end{aligned}$$

for positive integer  $l, 0 < l < k$ .

**Proof** Let  $v_k$  be the  $k$ -th derivative of  $u$ ; that is  $v_k := u^{(k)}$ , and denote  $v_0^k = u^k, v_0 = u, v_1 = u', v_2 = u'', v_1^2 = (u')^2$ , etc. Now let us use the mathematical induction to prove (6.2). When  $k = 1$ , we have

$$v_2 = \sum_{i=0}^{\lfloor \frac{C_{1,0}}{4} \rfloor} E_{1,i} u^{C_{1,i}} = E_{10} u^{C_{1,0}} = v_0^3$$

and

$$C_{0,0} = 1, \quad C_{1,0} = 3, \quad E_{1,0} = E_{0,0} C_{0,0} \left[ \frac{1}{2} (C_{0,0} - 1) + 1 \right] = 1.$$

Suppose that  $k \in \mathbb{N}$  and  $v_{2k} = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} \cdot v_0^{C_{k,i}}$ . Then by (6.1) we obtain

$$v_{2k+1} = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} C_{k,i} v_0^{C_{k,i}-1} v_1,$$

$$v_{2k+2} = \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} C_{k,i} v_0^{C_{k,i}-1} v_2 + \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} E_{k,i} C_{k,i} (C_{k,i} - 1) v_0^{C_{k,i}-2} v_1^2,$$

$$\begin{aligned}
 v_{2k+2} &= \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} O_{k,i} \left[ \frac{1}{2} (C_{k,i} + 1) \right] v_0^{C_{k,i}+2} + \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} O_{k,i} (C_{k,i} - 1) E(0) v_0^{C_{k,i}-2} \\
 &= \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} O_{k,i} \left[ \frac{1}{2} (C_{k,i} + 1) \right] v_0^{C_{k+1,i}} + \sum_{i=0}^{\lfloor \frac{C_{k,0}}{4} \rfloor} O_{k,i} (C_{k,i} - 1) E(0) v_0^{C_{k+1,i+1}} \\
 &= O_{k,0} \left[ \left( \frac{1}{2} (C_{k,0} + 1) \right) \right] v_0^{C_{k+1,0}} + O_{k,0} (C_{k,0} - 1) E(0) v_0^{C_{k+1,1}} \\
 &\quad + O_{k,1} \left[ \left( \frac{1}{2} (C_{k,1} + 1) \right) \right] v_0^{C_{k+1,1}} + O_{k,1} (C_{k,1} - 1) E(0) v_0^{C_{k+1,2}} \\
 &\quad + O_{k,2} \left[ \left( \frac{1}{2} (C_{k,2} + 1) \right) \right] v_0^{C_{k+1,2}} + \dots \\
 &\quad + O_{k, \lfloor \frac{C_{k,0}}{4} \rfloor} \left( C_{k, \lfloor \frac{C_{k,0}}{4} \rfloor} - 1 \right) E(0) v_0^{C_{k+1, \lfloor \frac{C_{k,0}}{4} \rfloor + 1}}.
 \end{aligned}$$

Hence

$$v_{2k+2} = \sum_{i=0}^{\lfloor \frac{C_{k+1,0}}{4} \rfloor} E_{k+1,i} \cdot v_0^{C_{k+1,i}},$$

which completes the induction, and we obtain (6.2). Using (6.2), we get (6.3).

### 7 The Blow-up Rate and Blow-up Constant for $u^{(k)}$

Finding out the blow-up rate and blow-up constant of  $u^{(k)}$  of (0.1) is our main result.

**Theorem 11** If  $u$  is the solution of problem (0.1) with one of the following properties that:

- (i)  $E(n_0) < 0$ ,
- (ii)  $E(n_0) = 0, a'(n_0) > 0$ ,
- (iii)  $E(n_0) > 0$ .

Then the blow-up rate of  $u^{(2k)}$  is  $1 + 2k$ , and the blow-up constant of  $u^{(2k)}$  is  $|E_{k,0}(\sqrt{2})^{1+2k}|$ ; that is, for  $k \in \mathbb{N}, m \in \{1, 2, 3, 4, 5, 6\}$ ,

$$\lim_{n \rightarrow N_m^*} u^{(2k)}(n) (N_m^* - n)^{1+2k} = (\pm 1)^{C_{k,0}} E_{k,0} (\sqrt{2})^{1+2k} := K_{2k}. \tag{7.1}$$

The blow-up rate of  $u^{(2k+1)}$  is  $2k+2$ , and the blow-up constant of  $u^{(2k+1)}$  is  $|E_{k,0}C_{k,0}(\sqrt{2})^{2k+1}|$ ; that is, for  $k \in \mathbb{N}, m \in \{1, 2, 3, 4, 5, 6\}$ ,

$$\lim_{n \rightarrow N_m^*} u^{(2k+1)}(n) (N_m^* - n)^{2+2k} = (\pm 1)^{C_{k,0}} E_{k,0}C_{k,0} (\sqrt{2})^{2k+1} := K_{2k+1}, \tag{7.2}$$

where  $C_{k,0} = 2k + 1, E_{k,0} = \prod_{i=0}^{k-1} [\frac{4i^2+5i+2}{2}]$ .

**Proof** Under condition (i),  $E(n_0) < 0, a'(n_0) \geq 0$  by (1.6) and (1.1), we get

$$\int_0^{J(n)} \frac{1}{N_1^* - n} \frac{dr}{\sqrt{\frac{1}{2} + E(n_0)r^4}} = 1, \quad \forall n \geq n_0. \tag{7.3}$$

Using Lemma 3 and (1.6), we obtain  $\lim_{n \rightarrow N_1^*} \sqrt{2} \frac{J(n)}{N_1^* - n} = 1$ , in other words,

$$\lim_{n \rightarrow N_1^*} a(n) (N_1^* - n)^2 = 2, \tag{7.4}$$

and then

$$\lim_{n \rightarrow N_1^*} u(n) (N_1^* - n) = \pm \sqrt{2}. \tag{7.5}$$

Here  $C_{k,i} = 2k + 1 - 4i$ , hence we have  $C_{k,i} > C_{k,j}$  as  $i < j$ . From (6.1) and (7.5), it follows

$$\lim_{n \rightarrow N_1^*} u^{(2k)}(n) (N_1^* - n)^{C_{k,0}} = (\pm 1)^{C_{k,0}} E_{k,0} (\sqrt{2})^{C_{k,0}}.$$

Since  $C_{k,0} = 1 + 2k$ , we get (7.1) for  $m = 1$ .

By (1.5), (7.4) and (6.2) we find that

$$\lim_{n \rightarrow N_1^*} J'(n) = -\frac{1}{\sqrt{2}}, \tag{7.6}$$

$$\begin{aligned} \sqrt{2} &= \lim_{n \rightarrow N_1^*} \left( a(n) (N_1^* - n)^2 \right)^{-\frac{3}{2}} \cdot \lim_{n \rightarrow N_1^*} a'(n) (N_1^* - n)^3, \\ \lim_{n \rightarrow N_1^*} u'(n) (N_1^* - n)^2 &= \pm \sqrt{2}, \end{aligned} \tag{7.7}$$

and

$$\begin{aligned} & \lim_{n \rightarrow N_1^*} u^{(2k+1)}(n) (N_1^* - n)^{C_{k,0}+1} \\ &= \lim_{n \rightarrow N_1^*} \sum_{i=0}^{k-1} E_{k,i} C_{k,i} u^{C_{k,i}-1}(n) \cdot u'(n) \cdot (N_1^* - n)^{C_{k,0}+1} \\ &= \lim_{n \rightarrow N_1^*} E_{k,0} C_{k,0} u^{C_{k,0}-1}(n) \cdot u'(n) \cdot (N_1^* - n)^{C_{k,0}+1} \\ &= \lim_{n \rightarrow N_1^*} E_{k,0} C_{k,0} u^{C_{k,0}-1}(n) \cdot (N_1^* - n)^{C_{k,0}-1} \cdot u' \cdot (N_1^* - n)^2 \\ &= (\pm)^{C_{k,0}} E_{k,0} C_{k,0} \left(\sqrt{2}\right)^{C_{k,0}}, \end{aligned}$$

thus (7.2) is proved for  $m = 1$ .

For  $E(0) < 0, a'(0) < 0$ , by (1.9) we have

$$\int_0^{J(n)} \frac{dr}{(N_2^* - n) \sqrt{\frac{1}{2} + E(n_0) r^4}} = 1 \quad \forall n \geq n_0. \tag{7.8}$$

Using Lemma 3, (7.8) and (6.1), therefore we gain the estimate (7.1) for  $m = 2$ , and by (1.8) we get estimate (7.2) for  $m = 2$ .

Under (ii),  $E(n_0) = 0, a'(n_0) > 0$ , we have

$$a(n) = a(n_0)^3 \left(\frac{1}{2} a'(n_0) (N_3^* - n)\right)^{-2} \quad \forall n \geq n_0. \tag{7.9}$$

In view of (7.9) and (6.1), we get estimate (7.1) for  $m = 3$ . Also, we have  $J'(n) = J'(n_0), \forall n \geq n_0$  and  $\lim_{n \rightarrow N_1^*} a(n)^{-\frac{3}{2}} a'(n) = -\frac{1}{2} a(n_0)^{-\frac{3}{2}} a'(n_0)$ .

By (7.9) and (6.2), estimate (7.2) for  $m = 3$  is completely proved.

Under (iii), the proofs of estimates (7.1) and (7.2) for  $m = 4, 5, 6$  are similar to the above, we omit the argumentation.

**Theorem 12** If  $u$  is the solution of problem (0.1) with  $E(n_0) > 0$  and  $a'(n_0) < 0$ , then we have

$$\lim_{n \rightarrow z^-(u_0, u_1)} u^{(2k)}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}} = (\pm)^{C_{k,k-1}} E_{k,k-1} E(n_0)^{\frac{C_{k,k-1}}{2}} \tag{7.10}$$

and

$$\lim_{n \rightarrow z^-(u_0, u_1)} u^{(2k+1)}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}+1} = E_{k,k-1} C_{k,k-1} E(n_0)^{C_{k,k-1}-1} \tag{7.11}$$

for  $k \in \mathbb{N}$ , where  $z$  is the null point (zero) of  $u$  and

$$C_{k,k-1} = 5 - 2n, \quad E_{k,k-1} = \prod_{i=0}^{k-1} (5 - 2i) (4 - 2i) E(n_0)^{k-1}.$$

**Proof** For  $E(n_0) > 0$  and  $a'(n_0) < 0$ , we have

$$\lim_{n \rightarrow z^-(u_0, u_1)} u^{(2k)}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}}$$

$$\begin{aligned}
&= \lim_{n \rightarrow z^-(u_0, u_1)} \sum_{i=0}^{k-1} E_{k,i} u^{C_{k,i}}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}} \\
&= \lim_{n \rightarrow z^-(u_0, u_1)} E_{k,k-1} u^{C_{k,k-1}}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}} \\
&= (\pm 1)^{C_{k,k-1}} E_{k,k-1} E(n_0)^{\frac{C_{k,k-1}}{2}}.
\end{aligned}$$

Therefore, (7.10) is proved.

From (6.2), we obtain that

$$\begin{aligned}
&\lim_{n \rightarrow z^-(u_0, u_1)} u^{(2k+1)}(n) (z(u_0, u_1) - n)^{-C_{k,k-1}+1} \\
&= \lim_{n \rightarrow z^-(u_0, u_1)} \sum_{i=0}^{k-1} E_{k,i} C_{k,i} u^{C_{k,i}-1}(n) u'(n) (z(u_0, u_1) - n)^{-C_{k,k-1}+1} \\
&= \lim_{n \rightarrow z^-(u_0, u_1)} E_{k,k-1} C_{k,k-1} u^{C_{k,k-1}-1}(n) u'(n) (z(u_0, u_1) - n)^{-C_{k,k-1}+1} \\
&= E_{k,k-1} C_{k,k-1} E(n_0)^{C_{k,k-1}}.
\end{aligned}$$

Thus, (7.11) is obtained.

**Acknowledgements** Thanks are due to Professor Tsai Long-Yi, Tai-Ping Liu, Professor Tong Yang and Professor Shih-Shien Yu for their continuous encouragement and discussions over this work, to Metta Education, Grand Hall and Auria Solar for their financial assistance.

### References

- [1] Li Mengrong. On the blow-up time and blow-up rate of positive solutions of semilinear wave equations  $\square u - u^p = 0$  in 1-dimensional space. *Commun Pure Appl Anal*, 2011
- [2] Li Mengrong. Estimates for the life-span of solutions of semilinear wave equations. *Commun Pure Appl Anal*, 2008, **7**(2): 417–432
- [3] Li Mengrong. On the semilinear wave equations. *Taiwanese J Math*, 1998, **2**(3): 329–345
- [4] Duan Renjun, Li Mengrong, Yang Tong. Propagation of singularities in the solutions to the Boltzmann equation near equilibrium. *Math Models Methods Appl Sci*, 2008, **18**(7): 1093–1114
- [5] Li Mengrong, Tsai Longyi. On a system of nonlinear wave equations. *Taiwanese J Math*, 2003, **7**(4): 557–573
- [6] Li Mengrong, Tsai Longyi. Existence and nonexistence of global solutions of some systems of semilinear wave equations. *Nonlinear Analysis*, 2003, **54**: 1397–1415
- [7] Li Mengrong, Pai Jente. Quenching problem in some semilinear wave equations. *Acta Math Sci*, 2008, **28B**(3): 523–529
- [8] Li Mengrong. On the generalized Emden-Fowler Equation  $u''(t)u(t) = c_1 + c_2u'(t)^2$  with  $c_1 \geq 0, c_2 \geq 0$ . *Acta Math Sci*, 2010, **30B**(4): 1227–1234
- [9] Shieh T H, Li M R. Numeric treatment of contact discontinuity with multi-gases. *J Comput Appl Math*, 2009, **230**(2): 656–673
- [10] Li M R, Lin Y J, Shieh T H. The flux model of the movement of tumor cells and health cells using a system of nonlinear heat equations. *Journal of Computational Biology*, to appear, 2011
- [11] Li M R, Shieh T H, Yue C J, Lee P, Li Y T. Parabola method in ordinary differential equation. *Taiwanese J Math*, to appear
- [12] Bellman R. *Stability Theory of Differential Equations*. Yew York: McGraw-Hill, 1953