

## BLOW-UP BEHAVIOR FOR A SEMILINEAR HEAT EQUATION WITH A NONLINEAR BOUNDARY CONDITION

SHENG-CHEN FU, JONG-SHENQ GUO AND JE-CHIANG TSAI

(Received December 28, 2001, revised September 9, 2002)

**Abstract.** We study the blow-up behaviors of solutions of a semilinear heat equation with a nonlinear boundary condition. Under certain conditions, we prove that the blow-up point occurs only at the boundary. Then, by applying the well-known method of Giga-Kohn, we derive the time asymptotic of solutions near the blow-up time. Finally, we prove that the blow-up is complete.

**1. Introduction.** In this paper, we study the following initial boundary value problem

$$(1.1) \quad u_t = u_{xx} + u^p, \quad x \in (0, 1), \quad t > 0,$$

$$(1.2) \quad u_x(0, t) = 0, \quad u_x(1, t) = u^q(1, t), \quad t > 0,$$

$$(1.3) \quad u(x, 0) = u_0(x), \quad x \in [0, 1],$$

where  $p, q$  are positive constants, and  $u_0(x)$  is a positive smooth function. For convenience, we always assume that

$$u'_0(0) = 0, \quad u'_0(1) = u_0^q(1).$$

We say that the solution  $u$  of the problem (1.1)–(1.3) blows up if there is a finite time  $T$  such that  $\max_{0 \leq x \leq 1} u(x, t) \rightarrow \infty$  as  $t \uparrow T$ . It has been shown in [11] that the solution  $u$  of the problem (1.1)–(1.3) blows up if and only if  $\max\{p, q\} > 1$ . In [11], they also studied the blow-up set and derived the upper and lower bounds for blow-up rate under certain conditions.

A point  $x_0$  is said to be a blow-up point for  $u$  if there is a sequence  $\{(x_n, t_n)\}$  such that  $x_n \rightarrow x_0$ ,  $t_n \rightarrow T$ , and  $u(x_n, t_n) \rightarrow \infty$  as  $n \rightarrow \infty$ . Under certain conditions, it is shown in [11] that the boundary point  $x = 1$  is the only blow-up point. This phenomenon of blow-up on the boundary has been observed and studied by many authors. We refer the readers to two nice survey papers [4] and [2] and the references cited therein. See also the references cited in [11].

We are concerned with the blow-up behaviors of solutions of the problem (1.1)–(1.3). Hence throughout this work we always assume that  $\max\{p, q\} > 1$ . In the sequel, we shall assume that the solution  $u$  of the problem (1.1)–(1.3) blows up at  $T < \infty$ . First, we study the blow-up set. We prove that blow-up point occurs only at the boundary point  $x = 1$ , if  $u'_0 \geq 0$  in  $[0, 1]$ . This improves the results of [11], where the monotonicity of  $u$  in time is

---

2000 *Mathematics Subject Classification.* Primary 35K60; Secondary 34A12, 35B40, 35K20.

This work was partly supported by the National Science Council of the Republic of China under the contract NSC 89-2115-M-003-014.

assumed. Furthermore, by deriving some a priori estimates, with the help of the lower and upper bounds for blow-up rate, we can apply the well-known Giga-Kohn transformation (cf. [8]) to derive the time asymptotic of solutions. This gives a more precise information of the blow-up behaviors. We remark here that a similar problem for the case when the heat operator is replaced by the porous medium operator in the half real line has been studied by de Pablo, Quirós and Rossi [3].

The next question is the possibility of continuation of solutions beyond the blow-up time. For more references on this subject, we refer the readers to the paper [5] and some references listed there. We show that the blow-up for the problem (1.1)–(1.3) is complete, i.e., solutions blowing up in finite time will be infinite identically after the blow-up time.

This paper is organized as follows. We study the blow-up set in Section 2. Some a priori estimates are given in Section 3. In Section 4, we study the self-similar solution for the critical case by an ordinary differential equation approach. Then we study the time asymptotic of the solution in Section 5. Finally, in Section 6 we prove that the blow-up is complete.

We thank the referee for helpful comments which improve some results in Section 2.

**2. Blow-up set.** Let  $u$  be the solution of the problem

$$(2.1) \quad u_t = u_{xx} + u^p, \quad x \in (0, 1), \quad 0 < t < T,$$

$$(2.2) \quad u_x(0, t) = 0, \quad u_x(1, t) = u^q(1, t), \quad 0 < t < T,$$

$$(2.3) \quad u(x, 0) = u_0(x), \quad x \in [0, 1],$$

where  $T$  is the blow-up time of  $u$ . Here as usual we always assume that  $u'_0(0) = 0$  and  $u'_0(1) = u_0^q(1)$ . We shall assume that  $u'_0 \geq 0$  in  $[0, 1]$ , so that by the maximum principle we have  $u_x > 0$  in  $(0, 1) \times (0, T)$ . We shall modify the method of Friedman and McLeod ([7]) to study the blow-up set.

**THEOREM 2.1.** *Suppose that  $p > 1$ . If  $u'_0 \geq 0$  in  $[0, 1]$ , then the blow-up occurs only at  $x = 1$ .*

**PROOF.** Suppose that there is a blow-up point  $a \in [0, 1)$ . Then there is a sequence  $\{(x_n, t_n)\}$  such that  $x_n \rightarrow a$ ,  $t_n \rightarrow T$ , and  $u(x_n, t_n) \rightarrow \infty$  as  $n \rightarrow \infty$ . Fix a constant  $d \in (a, 1)$ . By comparing the solution  $u$  with the function

$$u(x_n, t_n) \sin[\pi(x - d)/(1 - d)] \exp\{-[\pi/(1 - d)]^2(t - t_n)\}$$

for each  $n$  sufficiently large, it is easy to show that

$$u(x, t) \geq u(x_n, t_n) \sin[\pi(x - d)/(1 - d)] \exp\{-[\pi/(1 - d)]^2(t - t_n)\}$$

for  $x \in [d, 1]$  and  $t \geq t_n$ . Hence

$$(2.4) \quad \lim_{t \rightarrow T} u(x, t) = \infty$$

uniformly over  $x \in [b, c]$  for any compact subset  $[b, c]$  of  $(d, 1)$ .

Now, we fix a compact subset  $[b, c]$  of  $(d, 1)$ . We take any  $r \in (1, p)$  and consider the function

$$J(x, t) = u_x(x, t) - g(x)u^r(x, t),$$

where  $g(x) = \varepsilon \sin[\pi(x - b)/(c - b)]$  for some  $\varepsilon > 0$  to be determined. Using (2.4), there is a  $t_0 \in (0, T)$  such that

$$(2.5) \quad J_t - J_{xx} - (pu^{p-1} + 2rg'u^{r-1})J \geq 0$$

in  $[b, c] \times [t_0, T)$  for any  $\varepsilon > 0$ . By choosing  $\varepsilon > 0$  sufficiently small such that  $J(x, t_0) \geq 0$  for any  $x \in [b, c]$ , it follows from the maximum principle that  $J \geq 0$  in  $[b, c] \times [t_0, T)$ . Hence we have

$$(2.6) \quad u^{-r}(x, t)u_x(x, t) \geq g(x) \quad \text{in } [b, c] \times [t_0, T).$$

An integration of (2.6) leads to a contradiction. Hence the theorem follows. □

If  $p \leq 1$ , then  $q > 1$ , since  $\max\{p, q\} > 1$ .

LEMMA 2.2. *Let  $0 < p \leq 1$ . If  $u'_0 \geq 0$  in  $[0, 1]$ , then  $x = 1$  is the only blow-up point.*

PROOF. Since  $u'_0(1) > 0$ , there is a constant  $\delta \in (0, 1)$  such that  $u'_0 > 0$  in  $[1 - \delta, 1]$ . Set

$$\eta = \inf_{1-\delta \leq x \leq 1} \{u'_0(x)/u_0^q(x)\}, \quad M = (q - p) \sup_{[1-\delta, 1] \times [0, T)} u^{p-1}.$$

Then  $0 < \eta \leq 1$  and  $0 < M < \infty$ . Choose a positive integer  $n \geq 3$  such that  $n \geq M$  and a positive number  $\varepsilon < \min\{\eta, \delta^n\}$ . Define  $g(x) = (x - 1 + \varepsilon^{1/n})^n$  if  $1 - \varepsilon^{1/n} \leq x \leq 1$ ;  $g(x) = 0$ , otherwise. It is easy to see that  $g \in C^2([0, 1])$  and satisfies

$$(2.7) \quad 0 \leq g \leq \varepsilon, \quad g' \geq 0, \quad g'' \geq 0, \quad g'' \geq Mg.$$

Then, by using the fact  $q > 1$  and the maximum principle, it is easy to show that

$$g(x)u^q(x, t) \leq u_x(x, t)$$

for  $0 \leq x \leq 1$  and  $0 \leq t < T$ . Hence

$$(2.8) \quad u^{-q}(x, t)u_x(x, t) \geq g(x)$$

for  $0 \leq x \leq 1$  and  $0 \leq t < T$ . An integration of (2.8) shows that  $u$  cannot blow up at any point  $x < 1$ . This proves the lemma. □

Indeed, the condition  $u'_0 \geq 0$  in  $[0, 1]$  can be removed if  $0 < p \leq 1$ .

THEOREM 2.3. *If  $0 < p \leq 1$ , then  $x = 1$  is the only blow-up point.*

PROOF. We first extend the function  $u(x, t)$  to  $w(x, t)$  defined on  $[-1, 1] \times [0, T)$  so that  $w(x, t) = u(x, t)$  and  $w(-x, t) = w(x, t)$  for  $x \in [0, 1]$  and  $t \in [0, T)$ . Then  $w$  satisfies

$$\begin{aligned} w_t &= w_{xx} + w^p, & x \in (-1, 1), & \quad 0 < t < T, \\ w_x(-1, t) &= -w^q(-1, t), & w_x(1, t) &= w^q(1, t), & \quad 0 < t < T. \end{aligned}$$

Arguing as Lemma 1.2 of [9], there exists  $t^* \in (0, T)$  such that

$$n(t) := \#\{a \in [-1, 1] \mid w_x(a, t) = 0\}$$

is a constant for all  $t \geq t^*$ . Moreover, there are  $C^1$  functions  $s_0(t), \dots, s_{\pm l}(t)$ ,  $l \geq 0$ , from  $[t^*, T)$  to  $[-1, 1]$  such that

$$s_{-l}(t) < \dots < s_0(t) < \dots < s_l(t), \quad s_0(t) \equiv 0, \\ \{a \in [-1, 1] \mid w_x(a, t) = 0\} = \{s_{-l}(t), \dots, s_0(t), \dots, s_l(t)\} \quad \text{for } t \geq t^*,$$

and the limit  $s_i := \lim_{t \uparrow T} s_i(t)$  exists for all  $i$ . Since  $n(t)$  is constant in  $[t^*, T)$ , it follows from Theorem C of [1] that  $w_{xx}(s_i(t), t) \neq 0$  for all  $t \in [t^*, T)$ . Note that for each  $i$  there is a fixed sign for  $w_{xx}(s_i(t), t)$  for all  $t \in [t^*, T)$ . Also, it suffices to consider the so-called maximum curve, i.e., the curve for which  $w_{xx}(s_i(t), t) < 0$  on  $[t^*, T)$ .

If  $l = 0$ , then  $u_x(x, t) > 0$  on  $(0, 1] \times [t^*, T)$ . Hence the conclusion follows from Lemma 2.2. Suppose that  $l > 0$ . Set  $m_i(t) := w(s_i(t), t)$ . Notice that  $m'_i(t) < m_i(t)^p$  on  $[t^*, T)$ , if  $w_{xx}(s_i(t), t) < 0$  on  $[t^*, T)$ . Since  $0 < p \leq 1$ ,  $m_i(t)$  remains bounded near  $T$ . This implies that  $w$  cannot blow up at any point in  $(-1, 1)$ . The theorem is proved.  $\square$

**3. Some a priori estimates.** In this section, we will derive some a priori estimates which will be used in Section 5 to prove the time asymptotic results. Let  $u$  be the solution of (2.1)–(2.3) with blow-up time  $T$ . From now on we shall always assume that  $u'_0 \geq 0$  in  $[0, 1]$ , so that by the maximum principle we have  $u_x > 0$  in  $(0, 1) \times (0, T)$ . Notice that  $u(1, t) = \max_{0 \leq x \leq 1} u(x, t)$ .

The following lemma is given in [11] under the assumption  $u''_0 + u_0^p \geq a > 0$  in  $[0, 1]$ . Indeed, we have the following lemma.

LEMMA 3.1. *If  $u''_0 + u_0^p \geq 0$  in  $[0, 1]$ , then  $u_t \geq 0$  in  $[0, 1] \times [0, T)$ .*

PROOF. Set  $v = u_t$ . Then  $v$  satisfies

$$v_t = v_{xx} + pu^{p-1}v, \quad 0 < x < 1, \quad 0 < t < T, \\ v_x(0, t) = 0, \quad v_x(1, t) = qu^{q-1}(1, t)v(1, t), \quad 0 < t < T, \\ v(x, 0) = u''_0 + u_0^p \geq 0, \quad 0 \leq x \leq 1.$$

For any fixed  $\tau \in (0, T)$ , let

$$L = \max_{0 \leq x \leq 1, 0 \leq t \leq \tau} \left\{ \frac{1}{2}qu^{q-1}(x, t) \right\}, \quad M = 2L + 4L^2 + \max_{0 \leq x \leq 1, 0 \leq t \leq \tau} \{pu^{p-1}(x, t)\}.$$

Set  $w(x, t) = e^{-Mt-Lx^2}v(x, t)$ . Then  $w$  satisfies

$$w_t = w_{xx} + 4Lxw_x + cw, \quad 0 < x < 1, \quad 0 < t \leq \tau, \\ w_x(0, t) = 0, \quad w_x(1, t) = dw(1, t), \quad 0 < t \leq \tau, \\ w(x, 0) \geq 0, \quad 0 \leq x \leq 1,$$

where  $c = c(x, t) \leq 0$  and  $d = d(t) \leq 0$ . By the maximum principle, we obtain that  $w \geq 0$  in  $[0, 1] \times [0, \tau]$ . Hence the lemma follows.  $\square$

Recall from [11] that if  $u_t \geq 0$ , then there are positive constants  $c$  and  $A$  such that

$$(3.1) \quad c(T - t)^{-\alpha} \leq u(1, t) \leq A(T - t)^{-\alpha},$$

where the exponent is given by

$$\alpha = \begin{cases} 1/(p - 1) & \text{if } p \geq 2q - 1, \\ 1/[2(q - 1)] & \text{if } p < 2q - 1. \end{cases}$$

Hereafter we shall assume that  $u'_0 \geq 0$  and  $u''_0 + u^p_0 \geq 0$  in  $[0, 1]$ . Therefore, we have  $u_x \geq 0$  and  $u_t \geq 0$ . We now make the following Giga-Kohn transformation

$$(3.2) \quad y = \frac{1 - x}{\sqrt{T - t}}, \quad s = -\ln(T - t),$$

$$(3.3) \quad w(y, s) = (T - t)^\alpha u(x, t),$$

where  $\alpha$  is defined as in (3.1). Let

$$W = \{(y, s) \mid 0 < y < e^{s/2}, s > -\ln T\}.$$

Then for  $p > 2q - 1$  we have

$$(3.4) \quad w_s = w_{yy} - \frac{y}{2}w_y - \alpha w + w^p \quad \text{in } W,$$

$$(3.5) \quad w_y(0, s) = -e^{\gamma s}w(0, s)^q, \quad w_y(e^{s/2}, s) = 0, \quad s > -\ln T,$$

$$(3.6) \quad w(y, -\ln T) = T^\alpha u_0(1 - y\sqrt{T}), \quad 0 \leq y \leq 1/\sqrt{T},$$

where  $\gamma = [(2q - 1) - p]/[2(p - 1)] < 0$ ; for  $p = 2q - 1$  we have

$$(3.7) \quad w_s = w_{yy} - \frac{y}{2}w_y - \alpha w + w^p \quad \text{in } W,$$

$$(3.8) \quad w_y(0, s) = -w(0, s)^q, \quad w_y(e^{s/2}, s) = 0, \quad s > -\ln T,$$

$$(3.9) \quad w(y, -\ln T) = T^\alpha u_0(1 - y\sqrt{T}), \quad 0 \leq y \leq 1/\sqrt{T},$$

while for  $p < 2q - 1$  we have

$$(3.10) \quad w_s = w_{yy} - \frac{y}{2}w_y - \alpha w + e^{\sigma s}w^p \quad \text{in } W,$$

$$(3.11) \quad w_y(0, s) = -w(0, s)^q, \quad w_y(e^{s/2}, s) = 0, \quad s > -\ln T,$$

$$(3.12) \quad w(y, -\ln T) = T^\alpha u_0(1 - y\sqrt{T}), \quad 0 \leq y \leq 1/\sqrt{T},$$

where  $\sigma = [p - (2q - 1)]/[2(q - 1)] < 0$ .

We have the following a priori estimates for  $w$ .

LEMMA 3.2. *w and  $w_y$  are bounded in  $\bar{W}$ .*

PROOF. The fact that  $w$  is bounded follows from (3.1).

It follows from Lemma 3.1 that  $u_{xx} \geq -u^p$  in  $[0, 1] \times [0, T)$ . Multiplying the above inequality by  $u_x \geq 0$  and integrating it from  $x$  to 1, we obtain

$$(3.13) \quad u_x^2(x, t) \leq u^{2q}(1, t) + \frac{2}{p + 1}u^{p+1}(1, t).$$

Note that  $w_y(y, s) = -(T - t)^{\alpha+1/2}u_x(x, t)$ .

Recall (3.1). For  $p \geq 2q - 1$ , it follows from (3.13) and Lemma 3.1 that

$$\begin{aligned} w_y^2(y, s) &\leq (T - t)^{2\alpha+1}u^{2q}(1, t) + \frac{2}{p + 1}(T - t)^{2\alpha+1}u^{p+1}(1, t) \\ &= [(T - t)^\alpha u(1, t)]^{p+1}u^{2q-1-p}(1, t) + \frac{2}{p + 1}[(T - t)^\alpha u(1, t)]^{p+1} \\ &\leq A^{p+1}u_0^{2q-1-p}(1) + A^{p+1}. \end{aligned}$$

For  $p < 2q - 1$ , it also follows from (3.13) and Lemma 3.1 that

$$\begin{aligned} w_y^2(y, s) &\leq (T - t)^{2\alpha+1}u^{2q}(1, t) + \frac{2}{p + 1}(T - t)^{2\alpha+1}u^{p+1}(1, t) \\ &= [(T - t)^\alpha u(1, t)]^{2q} + \frac{2}{p + 1}[(T - t)^\alpha u(1, t)]^{2q}u^{p-2q+1}(1, t) \\ &\leq A^{2q} + \frac{2}{p + 1}A^{2q}u_0^{p-2q+1}(1). \end{aligned}$$

Hence the lemma is proved. □

LEMMA 3.3. *There is a positive constant  $C$  such that  $|w_s(y, s)| \leq C(1 + y)$  and  $|w_{yy}(y, s)| \leq C(1 + y)$  in  $\bar{W}$ .*

PROOF. It follows from Lemma 3.2 that  $|w_s(y, s) - w_{yy}(y, s)| \leq C(1 + y)$  in  $\bar{W}$  for some positive constant  $C$ . The lemma follows by applying the standard theory of parabolic equations, e.g., Theorem 6.44, Theorem 4.30 and Theorem 4.31 in [10]. □

**4. Self-similar solution.** In this section, we shall study the self-similar solution of (1.1)–(1.2) for the case  $p = 2q - 1$ , i.e.,  $q = (p + 1)/2$ . We are concerned with the existence and uniqueness of global positive monotone decreasing solution of the initial value problem (P):

$$(4.1) \quad w'' - \frac{1}{2}yw' - \alpha w + w^p = 0, \quad y > 0,$$

$$(4.2) \quad w'(0) = -w^q(0),$$

where  $w = w(y)$  and  $\alpha = 1/(p - 1)$ . We always assume that  $p > 1$ . The existence result has been obtained before by Wang and Wang in [12]. Here we present a different proof for the existence. Some of the lemmas will be useful for the proof of uniqueness.

Given any  $\eta > 0$ , there is a unique local solution  $w(y; \eta)$  of (4.1)–(4.2) with  $w(0) = \eta$ . Let  $\rho(y) = \exp\{-y^2/4\}$  and  $f(w) = -\alpha w + w^p$ . Then  $w$  satisfies

$$(4.3) \quad (\rho w')(y) = -\eta^q - \int_0^y \rho(s)f(w(s))ds.$$

Let  $\kappa$  be the unique positive solution of  $f(w) = 0$ . Note that  $w' < 0$  as long as  $w \geq \kappa$ . Set

$$(4.4) \quad \kappa_0 = \left(\frac{p+1}{p+3}\right)^\alpha \kappa.$$

Note that  $0 < \kappa_0 < \kappa$ .

LEMMA 4.1. *Let  $\eta \geq \kappa_0$ . Then  $w' < 0$  as long as  $w > 0$ .*

PROOF. Let

$$G(y) = \frac{1}{2}[w'(y)]^2 + F(w(y)),$$

where  $F(w) = \int_\kappa^w f(s)ds$ . Note that

$$(4.5) \quad G(0) \geq F(0) \quad \text{if and only if } \eta \geq \kappa_0.$$

Since

$$G'(y) = \frac{1}{2}y[w'(y)]^2,$$

and the problem (P) has no non-trivial constant solution,  $G$  is strictly increasing.

If  $w$  is not monotone decreasing, then there is the first critical point  $y_0 > 0$  of  $w$  such that  $w'(y_0) = 0$  and  $w > 0$  in  $[0, y_0]$ . Notice that  $w(y_0) < \kappa$ . Hence

$$G(y_0) = F(w(y_0)) < F(0).$$

On the other hand, by (4.5) we have  $G(0) \geq F(0)$ , since  $\eta \geq \kappa_0$ . This implies that  $G(0) > G(y_0)$ , a contradiction. Therefore, the lemma follows.  $\square$

Suppose that  $w > 0$  and  $w' < 0$  in  $[0, \infty)$ . Let  $l = \lim_{y \rightarrow \infty} w(y)$ . Then  $l \in [0, \kappa)$  and there is a sequence  $\{y_n\}$  such that  $w'(y_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Dividing Equation (4.1) by  $y$  and integrating it from 1 to  $y_n$  for any  $n$  large, as  $n \rightarrow \infty$ , this leads to a contradiction, if  $l \in (0, \kappa)$ . Hence  $l = 0$ .

LEMMA 4.2. *For  $\eta \geq \kappa_0$ , the solution  $w$  is monotone decreasing to zero at some finite  $R$ .*

PROOF. Otherwise, by Lemma 4.1 and the above observation,  $w(y) \rightarrow 0$  as  $y \rightarrow \infty$  and there is a sequence  $\{y_n\}$  such that  $w'(y_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then  $G(y_n) \rightarrow F(0)$  as  $n \rightarrow \infty$ . Since  $G$  is monotone increasing, its limit must be greater than  $G(0)$ , i.e.,  $F(0) > G(0)$ , a contradiction to (4.5). This proves the lemma.  $\square$

We now turn to the case when  $\eta$  is small. First, let  $\eta_0$  be a positive constant such that  $-f(w) \geq \alpha w/2$  for all  $w \in [0, \eta_0]$ . Notice that  $\eta_0 < \kappa$ . Choose  $\eta_1 \in (0, \eta_0)$  such that  $\eta^{1-q} > e^{1/4}$  for all  $\eta \in (0, \eta_1)$ . Now, given any fixed  $\eta \in (0, \eta_1)$ , suppose that  $w' < 0$  in  $[0, R]$  and  $w(R) = 0$  for some  $R = R(\eta) > 0$ . Since, by (4.3),  $\rho(y)w'(y) \geq -\eta^q$  for all  $y \in [0, R)$ , we have

$$\eta = - \int_0^R w'(s)ds \leq \eta^q R e^{R^2/4}.$$

Let  $g(y) = ye^{y^2/4}$ . Since  $g$  is strictly monotone increasing, we conclude that  $R = R(\eta) > 1$ , if  $\eta < \eta_1$ .

LEMMA 4.3. *There is a small positive constant  $\eta_*$  such that  $w'$  vanishes before  $w$  vanishes, if  $\eta < \eta_*$ .*

PROOF. Let  $\eta_*$  be a positive constant such that  $\eta_* < \eta_1$  and

$$(4.6) \quad \eta^{1-q} > \frac{1 + \alpha/4}{\alpha/2} e^{1/4} \quad \text{for all } \eta \in (0, \eta_*).$$

Suppose that there is an  $\eta \in (0, \eta_*)$  such that the lemma does not hold. Then the corresponding solution  $w$  must have the property that  $w' < 0$  in  $[0, y_0]$  for some  $y_0 > 1$ . From (4.3) it follows that  $\rho(y)w'(y) \geq -\eta^q$  for all  $y \in [0, y_0]$  and so

$$(4.7) \quad w(y) = \eta + \int_0^y w'(s)ds \geq \eta - \eta^q ye^{y^2/4} \quad \text{for all } y \in [0, y_0].$$

Then from (4.3), (4.7), and noting that  $\eta < \eta_0$ , we obtain that

$$\begin{aligned} \rho(y)w'(y) &\geq -\eta^q + \frac{\alpha}{2} \int_0^y \rho(s)w(s)ds \\ &\geq -\eta^q + \frac{\alpha}{2} \int_0^y e^{-s^2/4} [\eta - \eta^q se^{s^2/4}] ds \\ &\geq -\left(1 + \frac{\alpha}{4}y^2\right)\eta^q + \frac{\alpha}{2}ye^{-y^2/4}\eta \end{aligned}$$

for all  $y \in (0, y_0)$ . In particular, for  $y = 1$  we have

$$e^{-1/4}w'(1) \geq -\left(1 + \frac{\alpha}{4}\right)\eta^q + \frac{\alpha}{2}e^{-1/4}\eta > 0,$$

since  $\eta < \eta_*$ . This is a contradiction and the lemma is proved. □

Now, we define

$$\begin{aligned} I_1 &= \{\eta > 0 \mid w(y; \eta) \text{ is decreasing to zero at some finite } R\} \\ I_2 &= \{\eta > 0 \mid w'(y; \eta) \text{ vanishes before } w(y; \eta) \text{ vanishes}\} \end{aligned}$$

Notice that  $w$  and  $w'$  cannot vanish at the same time. Hence  $I_1$  and  $I_2$  are disjoint. Lemmas 4.2 and 4.3 imply that  $[\kappa_0, \infty) \subset I_1$  and  $(0, \eta_*) \subset I_2$ .

LEMMA 4.4. *The set  $I_2$  is open.*

PROOF. Let  $\eta_0 \in I_2$ . Then  $\eta_0 < \kappa_0 < \kappa$  and there is the first point  $y_0 > 0$  such that  $w_0 > 0$  in  $[0, y_0]$ ,  $w'_0 < 0$  in  $[0, y_0)$  and  $w'_0(y_0) = 0$ , where  $w_0(y) = w(y; \eta_0)$ . Since  $w''_0(y_0) > 0$ , there is a positive constant  $\delta$  such that  $w'_0(y) > 0$  for  $y \in (y_0, y_0 + \delta]$ . Let  $\varepsilon > 0$ ,  $\varepsilon < w_0(y_0)/2$ , and  $\varepsilon < w'_0(y_0 + \delta)/2$ . By the continuous dependence of initial value, there is a positive constant  $\gamma$  such that  $|w(y; \eta) - w_0(y)| < \varepsilon$  and  $|w'(y; \eta) - w'_0(y)| < \varepsilon$  for all  $y \in [0, y_0 + \delta]$ , if  $\eta \in (\eta_0 - \gamma, \eta_0 + \gamma)$ . This implies that  $(\eta_0 - \gamma, \eta_0 + \gamma) \subset I_2$  and so  $I_2$  is open. □



To prove that  $I_1$  is open, we consider the quantity

$$(4.8) \quad H(y) = \alpha w(y) + \frac{1}{2}yw'(y).$$

Then  $H$  satisfies the equation

$$(4.9) \quad H'(y) = \frac{1}{2}yH(y) + \left(\frac{1}{2} + \alpha\right)w'(y) - \frac{1}{2}yw^p(y).$$

Suppose that  $H(y_0) < 0$  for some  $y_0 \geq 0$ . Then  $w'(y_0) < 0$  by (4.8) and  $H'(y_0) < 0$  by (4.9). Hence, by (4.9) again,  $H'(y) < 0$  and  $H(y) < 0$  for all  $y \geq y_0$  as long as  $w > 0$ .

LEMMA 4.5. *The set  $I_1$  is open.*

PROOF. First, we claim that if there is a point  $y_0 \geq 0$  such that  $H(y_0) < 0$ , then  $w$  is decreasing after  $y_0$  and vanishes at some finite  $R > y_0$ . Otherwise, if  $w > 0$  in  $[0, \infty)$ , then  $H(y) < 0$  and  $H'(y) < 0$  for all  $y \geq y_0$ . Hence there is a positive constant  $\delta$  such that  $H(y) \leq -\delta$  for all  $y \geq y_0$ . By an integration, we obtain that

$$w(y) \leq (y_0/y)^{2\alpha}w(y_0) - \frac{\delta}{\alpha} + \frac{\delta}{\alpha}(y_0/y)^{2\alpha} \rightarrow -\frac{\delta}{\alpha} \quad \text{as } y \rightarrow \infty,$$

a contradiction.

Now, let  $\eta_0 \in I_1$  and  $w_0(y) = w(y; \eta_0)$ . Then there is a finite  $R_0 > 0$  such that  $w'_0 < 0$  and  $w_0 > 0$  in  $[0, R_0)$ . Since  $w_0(R_0) = 0$  and  $w'_0(R_0) < 0$ , there is a positive constant  $\delta$  such that  $H_0(R_0 - \delta) < 0$ , where  $H_0(y) = \alpha w_0(y) + yw'_0(y)/2$ . It follows from the theory of continuous dependence on initial value that there is a positive constant  $\gamma$  such that  $w(y; \eta) > 0$ ,  $w'(y; \eta) < 0$  for  $y \in [0, R_0 - \delta]$ , and  $H(R_0 - \delta) < 0$ , if  $\eta \in (\eta_0 - \gamma, \eta_0 + \gamma)$ . Then  $w$  is decreasing after  $R_0 - \delta$  and vanishes at some finite  $R > R_0 - \delta$ , if  $\eta \in (\eta_0 - \gamma, \eta_0 + \gamma)$ . Hence the lemma is proved.  $\square$

We now state and prove an existence theorem as follows.

THEOREM 4.6. *There is a global positive monotone decreasing solution of (P).*

PROOF. Set  $\bar{\eta} = \inf I_1$ . Then the corresponding solution  $\bar{w}(y) = w(y; \bar{\eta})$  must be a global positive monotone decreasing solution of (P).  $\square$

Indeed, for any  $\eta \notin I_1 \cup I_2$ , the corresponding solution  $w(y; \eta)$  is a global positive monotone decreasing solution of (P) satisfying  $w(y; \eta) \rightarrow 0$  as  $y \rightarrow \infty$ .

We have from Lemma 4.2 the estimate  $\bar{\eta} < \kappa_0$ . Also, the initial value  $\eta < \kappa_0$  for any global positive monotone decreasing solution of (P). To derive a better estimate, we need the following generalized version of Pohozaev Identity, which is inspired by Lemma 2.1 of [13] (see also [14]).

LEMMA 4.7. *Suppose  $w(y)$  is a solution of (P) and define*

$$J(y) := \rho(y)(w'(y))^2 - \frac{y}{2}\rho(y)w'(y)w(y) + \left(\frac{1}{4} - \alpha\right)\rho(y)w^2(y) + \frac{2}{p+1}\rho(y)w^{p+1}(y),$$

where  $\rho(y) = \exp[-y^2/4]$ . Then the following identity holds:

$$J(y) = \frac{p+3}{p+1}w^{p+1}(0) + \frac{p-5}{4(p-1)}w^2(0) + \int_0^y s\rho(s) \left\{ \frac{p-1}{2(p+1)}w^{p-1}(s) - \frac{1}{8} \right\} w^2(s)ds.$$

PROOF. Differentiating  $J(y)$  and using (4.1), we obtain

$$J'(y) = y\rho(y) \left\{ \frac{p-1}{2(p+1)}w^{p-1}(y) - \frac{1}{8} \right\} w^2(y).$$

Integrating  $J'(y)$  from 0 to  $y$  and noting that

$$J(0) = \frac{p+3}{p+1}w^{p+1}(0) + \frac{p-5}{4(p-1)}w^2(0),$$

we get the desired identity. □

COROLLARY 4.8. Suppose that  $w(y)$  is a global positive solution of (P) satisfying  $w(y) \rightarrow 0$  as  $y \rightarrow \infty$ . Then

$$(4.10) \quad \int_0^\infty s\rho(s) \left\{ \frac{1}{8} - \frac{p-1}{2(p+1)}w^{p-1}(s) \right\} w^2(s)ds = \frac{p+3}{p+1}w^{p+1}(0) + \frac{p-5}{4(p-1)}w^2(0).$$

PROOF. Since  $w(y)$  is a global positive solution of (P) and  $\lim_{y \rightarrow \infty} w(y) = 0$ , there is a sequence  $y_n \rightarrow \infty$  such that  $\lim_{n \rightarrow \infty} w'(y_n) = 0$ . Using Lemma 4.7, (4.10) follows. □

Define  $f_1(w) = \{1/8 - [(p-1)/(2(p+1))]w^{p-1}\}w^2$  and let  $\bar{\kappa} = (\alpha/2)^\alpha$ . Then it is easy to check that

$$\max_{w \in [0, \infty)} f_1(w) = f_1(\bar{\kappa}).$$

The following lemma gives an upper bound for any global positive solution of (P) which tends to zero as  $y \rightarrow \infty$ .

LEMMA 4.9. Suppose that  $w(y)$  is a global positive solution of (P) with  $w(0) = \eta$  such that  $w(y) \rightarrow 0$  as  $y \rightarrow \infty$ . Then  $\eta < \bar{\kappa}$ . In particular, we have  $\bar{\eta} < \bar{\kappa}$ .

PROOF. For contradiction, we assume that  $\eta \geq \bar{\kappa}$ . It follows from the definition of  $\bar{\kappa}$  that

$$\int_0^\infty s\rho(s)f_1(w(s))ds < \int_0^\infty s\rho(s)f_1(\bar{\kappa})ds = \frac{p-1}{4(p+1)}\bar{\kappa}^2.$$

On the other hand, since  $w(0) = \eta \geq \bar{\kappa}$ , we have

$$\begin{aligned} \frac{p+3}{p+1}w^{p+1}(0) + \frac{p-5}{4(p-1)}w^2(0) &\geq \frac{p+3}{2(p+1)(p-1)}w^2(0) + \frac{p-5}{4(p-1)}w^2(0) \\ &= \frac{p-1}{4(p+1)}w^2(0) \geq \frac{p-1}{4(p+1)}\bar{\kappa}^2, \end{aligned}$$

a contradiction to (4.10). This completes the proof. □

THEOREM 4.10. If  $1 < p \leq 2$ , then there is a unique global positive monotone decreasing solution of (P).

PROOF. For contradiction, we suppose that there are two distinct global positive monotone decreasing solutions  $w_1$  and  $w_2$  of (P). Note that  $w_i(y) \rightarrow 0$  as  $y \rightarrow \infty$  for  $i = 1, 2$ .

First, we claim that  $w_1$  and  $w_2$  must intersect each other at least once. Multiplying the equation

$$(\rho w_i')'(y) = -\rho(y)f(w_i(y)), \quad i = 1, 2,$$

by  $w_2$  for  $i = 1$ ; by  $w_1$  for  $i = 2$ , respectively, and integrating by parts, we end up with

$$\int_0^\infty \rho(s)w_1(s)w_2(s)(w_1^{p-1}(s) - w_2^{p-1}(s))ds = w_1(0)w_2(0)(w_2^{q-1}(0) - w_1^{q-1}(0)),$$

using  $w_i'(0) = -w_i^q(0)$ ,  $i = 1, 2$ . Hence they must intersect each other at least once.

Without loss of generality, we may assume that  $w_1(y) > w_2(y)$  in  $[0, y_0)$  and  $w_1(y_0) = w_2(y_0)$  for some  $y_0 > 0$ . Then we have  $w_1'(y_0) < w_2'(y_0)$ . Hence there exists  $y_1 > y_0$  such that  $w_1(y_1) < w_2(y_1)$ . Define  $v(y) := w_1(y) - w_2(y)$ . Then it follows from (4.1) that  $v(y)$  satisfies

$$(4.11) \quad v'' - \frac{y}{2}v' + [p\xi^{p-1}(y) - \alpha]v = 0,$$

for some  $\xi(y) \in [\min\{w_1(y), w_2(y)\}, \max\{w_1(y), w_2(y)\}]$ . Since  $\lim_{y \rightarrow \infty} w_i(y) = 0$ ,  $i = 1, 2$ , there exists  $y_2 > y_1$  such that  $|v(y_2)| < |v(y_1)|/2$ .

Now, let  $\bar{y} \in [0, y_2]$  be a minimal point of  $v$  in  $[0, y_2]$ . Then  $\bar{y} \in (0, y_2)$  and  $v(\bar{y}) < 0$ . Since  $\bar{y}$  is an interior extreme point of  $v$ , we have

$$(4.12) \quad v'(\bar{y}) = 0, \quad v''(\bar{y}) \geq 0.$$

From Lemma 4.9 and  $\xi(\bar{y}) \leq \max\{w_1(0), w_2(0)\}$ , it follows that

$$(4.13) \quad p\xi^{p-1}(\bar{y}) - \alpha < 0,$$

if  $1 < p \leq 2$ . Then, by (4.11), (4.12) and (4.13), we obtain that

$$\begin{aligned} 0 &= v''(\bar{y}) - \frac{\bar{y}}{2}v'(\bar{y}) + [p\xi^{p-1}(\bar{y}) - \alpha]v(\bar{y}) \\ &\geq [p\xi^{p-1}(\bar{y}) - \alpha]v(\bar{y}) \\ &> 0, \end{aligned}$$

a contradiction. This completes the proof. □

We conjecture that Theorem 4.10 should hold for any  $p > 1$ . Unfortunately we are unable to prove it now, so we left it as an open problem.

**5. Time asymptotic analysis.** In this section, we shall study the time asymptotic of the solutions of the problem (2.1)–(2.3) for various cases. The method is the same as the one used in [8] with some modifications. Hence we shall only give the outline of the proofs.

**THEOREM 5.1.** *For  $p > 2q - 1$ , we have*

$$(T - t)^\alpha u(1 - y\sqrt{T - t}, t) \rightarrow \kappa$$

as  $t \rightarrow T$  uniformly for  $y \in [0, C]$  for any  $C > 0$ . Here  $\alpha = 1/(p - 1)$  and  $\kappa = \alpha^\alpha$ .

PROOF. As in [8], we take any increasing sequence  $\{s_j\}$  in  $(0, \infty)$  such that  $s_{j+1} - s_j \rightarrow \infty$  as  $j \rightarrow \infty$ . For each  $j \in \mathbf{N}$ , we define

$$w_j(y, s) = w(y, s + s_j) \quad \text{for all } (y, s) \in W_j \equiv \{(y, s) \mid 0 \leq y \leq e^{(s+s_j)/2}, s \geq -s_j - \ln T\}.$$

Note that  $\bigcup_{j=1}^\infty W_j = [0, \infty) \times \mathbf{R}$  and  $W_1 \subset W_2 \subset \dots$ . Recall Lemmas 3.2 and 3.3. By the Ascoli-Arzelà Theorem and a diagonal process, we can get a subsequence (still denoted by  $w_j$ ) such that  $w_j(y, s) \rightarrow w_\infty(y, s)$  as  $j \rightarrow \infty$  uniformly on any compact subset of  $[0, \infty) \times \mathbf{R}$  and that for any integer  $m$  we have  $w_{j,y}(y, m) \rightarrow w_{\infty,y}(y, m)$  as  $j \rightarrow \infty$  pointwise for  $y \in [0, \infty)$  for some function  $w_\infty$  defined on  $[0, \infty) \times \mathbf{R}$ . It is easy to see that  $w_\infty$  is a classical solution of the equation

$$w_s = w_{yy} - \frac{1}{2}yw_y - \alpha w + w^p \quad \text{in } [0, \infty) \times \mathbf{R}.$$

Now, we claim that  $w_{\infty,s}(y, s) \equiv 0$  in  $[0, \infty) \times \mathbf{R}$ . Introduce the energy function

$$E[w](s) = \frac{1}{2} \int_0^s \rho w_y^2 dy + \frac{\alpha}{2} \int_0^s \rho w^2 dy - \frac{1}{p+1} \int_0^s \rho w^{p+1} dy,$$

where  $\rho(y) = e^{-y^2/4}$ . By a simple computation, we get

$$(5.1) \quad -\frac{d}{ds}E[w](s) = \int_0^s \rho w_s^2 dy - G(s),$$

where

$$G(s) = \rho(s) \left\{ \frac{1}{2}w_y^2(s, s) + \frac{\alpha}{2}w^2(s, s) - \frac{1}{p+1}w^{p+1}(s, s) + w_y(s, s)w_s(s, s) \right\} \\ + \exp \left\{ \frac{(2q-1)-p}{2(p-1)}s \right\} w(0, s)^q w_s(0, s).$$

Let  $s_0 = \max\{2 \ln 2, -\ln T\}$ . Note that

$$\{(y, s) \mid 0 \leq y \leq s, s \geq s_0\} \subseteq \bar{W}.$$

Integrating both sides of (5.1) from  $m + s_j$  to  $m + s_{j+1}$  for any  $m, j \in \mathbf{Z}$  with  $m + s_j \geq s_0$ , we obtain

$$\int_{m+s_j}^{m+s_{j+1}} \int_0^s \rho(y)w_s^2(y, s)dyds \\ = E_{m+s_j}[w](m + s_j) - E_{m+s_{j+1}}[w](m + s_{j+1}) + \int_{m+s_j}^{m+s_{j+1}} G(s)ds.$$

By a change of variable, we get

$$\int_m^{m+s_{j+1}-s_j} \int_0^{s+s_j} \rho(y)w_{j,s}^2(y, s)dyds \\ = E_{m+s_j}[w_j](m) - E_{m+s_{j+1}}[w_{j+1}](m) + \int_{m+s_j}^{m+s_{j+1}} G(s)ds.$$

Since

$$|G(s)| \leq C \exp \left\{ \frac{(2q-1)-p}{2(p-1)} s \right\} (1+s),$$

it follows that

$$\int_{s_0}^{\infty} |G(s)| ds < \infty.$$

Proceeding as in [8], we get

$$\int_m^M \int_0^{\infty} \rho w_{\infty,s}^2 dy ds = 0 \quad \text{for all } m, M \in \mathbf{Z}, \quad m < M.$$

Hence  $w_{\infty,s} \equiv 0$  and so  $w_{\infty}(y, s) = w_{\infty}(y)$  for all  $y \in [0, \infty)$  and  $s$ .

Note that  $w_{\infty}(0) > 0$ . Also, from  $w_{j,y}(0, s) = -e^{\gamma(s+s_j)} w_j(0, s)^q$ , where

$$\gamma = [(2q-1)-p]/[2(p-1)] < 0,$$

it follows that  $w'_{\infty}(0) = 0$ . Therefore,  $w_{\infty}$  is a bounded positive global solution of

$$w'' - \frac{1}{2}yw' - \alpha w + w^p = 0$$

and so  $w_{\infty} \equiv \kappa$  (cf. [8]). Since the sequence  $\{s_j\}$  is arbitrary, the theorem follows. □

Recall from [6] that there is a unique bounded positive global solution (denoted by  $V(y)$ ) of

$$(5.2) \quad w'' - \frac{1}{2}yw' - \alpha w = 0, \quad w'(0) = -w^q(0).$$

THEOREM 5.2. For  $p < 2q - 1$ , we have

$$(T-t)^{\alpha} u(1-y\sqrt{T-t}, t) \rightarrow V(y)$$

as  $t \rightarrow T$  uniformly for  $y \in [0, C]$  for any  $C > 0$ . Here  $\alpha = 1/[2(q-1)]$ .

PROOF. Let  $s_j, w_j, w_{\infty}$  be defined as in Theorem 5.1. Then it is easy to see that  $w_{\infty}$  is a classical solution of

$$w_s = w_{yy} - \frac{1}{2}yw_y - \alpha w, \quad y \in [0, \infty), \quad s \in \mathbf{R}.$$

Next, we introduce the energy function

$$E[w](s) = \frac{1}{2} \int_0^s \rho w_y^2 dy + \frac{\alpha}{2} \int_0^s \rho w^2 dy - \frac{1}{q+1} w^{q+1}(0, s).$$

Proceeding as in the proof of Theorem 5.1, we obtain that  $w_{\infty,s} \equiv 0$  and so  $w_{\infty}(y, s) = w_{\infty}(y)$ . Since  $w_y(0, s) = -w^q(0, s)$ , we get  $w'_{\infty}(0) = -w^q_{\infty}(0)$ . Recall  $w_{\infty}(0) > 0$ . Hence  $w_{\infty}(y) = V(y)$  and the theorem follows. □

Finally, we shall consider the critical case, i.e., the case  $p = 2q - 1$ . Suppose that  $\bar{w}(y)$  (as defined in Section 4) is the unique global positive monotone decreasing solution of (4.1). Then the same argument as above leads to the following conclusion. Note that  $w_y < 0$  for  $y \geq 0$ . Hence the limit function satisfies  $w'_{\infty} \leq 0$ .

THEOREM 5.3. *Let  $p = 2q - 1$ . If  $1 < p \leq 2$ , then we have*

$$(T - t)^\alpha u(1 - y\sqrt{T - t}, t) \rightarrow \bar{w}(y)$$

as  $t \rightarrow T$  uniformly for  $y \in [0, C]$  for any  $C > 0$ . Here  $\alpha = 1/(p - 1)$ .

**6. Complete blow-up.** Suppose that the solution  $u$  of the problem (1.1)–(1.3) blows up at the finite time  $T$  and  $x = 1$  is the only blow-up point. This is the case if  $\max\{p, q\} > 1$  and  $u'_0 \geq 0$  in  $[0, 1]$ . Let

$$f^{(n)}(s) = \min\{s^q, n^q\}, \quad g^{(n)}(s) = \min\{s^p, n^p\}, \quad s \geq 0, \quad n \in \mathbb{N},$$

and let  $u^{(n)}$  be the solution of the problem  $(B^{(n)})$ :

$$(6.1) \quad u_t^{(n)} = u_{xx}^{(n)} + g^{(n)}(u^{(n)}), \quad x \in (0, 1), \quad t > 0,$$

$$(6.2) \quad u_x^{(n)}(0, t) = 0, \quad u_x^{(n)}(1, t) = f^{(n)}(u^{(n)}(1, t)), \quad t > 0,$$

$$(6.3) \quad u^{(n)}(x, 0) = u_0(x), \quad x \in [0, 1].$$

We shall follow the method used in [5] to prove that the blow-up is complete, i.e., as  $n \rightarrow \infty$ ,  $u^{(n)}(x, t) \rightarrow \infty$  for all  $(x, t) \in [0, 1] \times (T, \infty)$ .

Let  $K = \max_{0 \leq x \leq 1} u_0(x)$ . Since  $f^{(n)}$  and  $g^{(n)}$  are locally Lipschitz in  $(0, K]$  and  $u'_0(1) = f^{(n)}(u_0(1))$  for  $n > K$ , the solution  $u^{(n)}$  of  $(B^{(n)})$  is  $C^1$  up to the boundary.

Suppose that  $v^{(n)}$  is a positive smooth supersolution and  $w^{(n)}$  is a smooth subsolution of  $(B^{(n)})$ . Then it is easy to show by the maximum principle that  $v^{(n)} \geq w^{(n)}$  for  $0 \leq x \leq 1$ ,  $t > 0$ , if  $n > K$ . Note that the function  $K + (n^p + n^q)t + n^q x^2/2$  is a supersolution of  $(B^{(n)})$ . Therefore, for any positive integer  $n > K$ , the problem  $(B^{(n)})$  has a unique positive global (in time) solution  $u^{(n)}$  such that  $u^{(n)} \leq u^{(n+1)}$  for  $(x, t) \in [0, 1] \times [0, \infty)$  and  $u^{(n)} \leq u$  for  $(x, t) \in [0, 1] \times [0, T)$ .

Now, we define

$$v(x, t) = \lim_{n \rightarrow \infty} u^{(n)}(x, t), \quad 0 \leq x \leq 1, \quad t > 0.$$

Then we can show that  $v(x, t) = u(x, t)$  for  $0 \leq t < T$ . Note that  $v(1, T) = \infty$ . Furthermore, we have

LEMMA 6.1. *If  $q \geq 1$ , then  $v(1, t) = \infty$  for all  $t \geq T$ .*

PROOF. For any  $M > 1$ , there is a smooth function  $U$  such that  $U(1) = M$ ,  $U'(1) = M^q$ ,  $U(\xi) = 0$ , and  $U'' + U^p = 0$  in  $(\xi, 1]$  for some  $\xi \in (0, 1)$ , since  $q \geq 1$ . We extend the function  $U$  to be linear on  $[0, \xi]$  so that  $U \in C^2([0, 1])$ . Let  $M > \max\{1, u_0(1), \|u'_0\|_\infty^{1/q}\}$ . Then  $u_0$  intersects  $U$  exactly once.

Since  $v(1, T) = \infty$ , there is a positive integer  $k > K$  such that  $u^{(k)}(1, t_0) > M$  for some  $t_0 \in (0, T)$ . Then there is  $t_1 \in (0, t_0)$  such that  $u^{(k)}(1, t_1) = M$  and  $u^{(k)}(1, t) < M$  for all  $t \in [0, t_1)$ . Since  $u^{(k)}(0, t) > U(0)$  and  $u^{(k)}(1, t) < U(1)$  for all  $t \in [0, t_1)$ , it implies that  $u^{(k)}(\cdot, t)$  intersects  $U$  at least once. Note that

$$(u^{(k)} - U)_t = (u^{(k)} - U)_{xx} + c(x, t)(u^{(k)} - U), \quad c(x, t) = \frac{g^{(k)}(u^{(k)}) + U_{xx}}{u^{(k)} - U}.$$

Since  $u^{(k)}(x, t)$  is bounded away from zero in  $[0, 1] \times [0, t_1]$ , the function  $c(x, t)$  is bounded. Applying Theorem D of [1],  $u^{(k)}(\cdot, t)$  intersects  $U$  exactly once for  $t < t_1$ . Let  $s^{(k)}(t)$  be the function such that

$$(u^{(k)} - U)(s^{(k)}(t), t) = 0 \quad \text{for all } t \in [0, t_1].$$

Applying Theorem D of [1] again, we have

$$(u^{(k)} - U)_x(s^{(k)}(t), t) \neq 0 \quad \text{for all } t \in [0, t_1].$$

Therefore, by the Implicit Function Theorem, the function  $s^{(k)}(t)$  is continuous in  $[0, t_1]$ .

Now, we will show that

$$(6.4) \quad u^{(k)}(x, t_1) \geq U(x) \quad \text{for all } x \in [0, 1].$$

To prove (6.4), we consider two cases. First, we suppose that  $\lim_{t \rightarrow t_1^-} s^{(k)}(t)$  exists. we claim that  $\lim_{t \rightarrow t_1^-} s^{(k)}(t) = 1$ . For contradiction, we assume that  $0 \leq \lim_{t \rightarrow t_1^-} s^{(k)}(t) < 1$ . Recall that  $(u^{(k)} - U)(1, t_1) = 0$  and  $(u^{(k)} - U)(x, t) \leq 0$  for all  $x \in [s^{(k)}(t), 1]$  and  $t \in [0, t_1]$ . By the Hopf Boundary Point Lemma,  $(u^{(k)} - U)_x(1, t_1) > 0$ , a contradiction. Since

$$(u^{(k)} - U)(x, t) \geq 0, \quad x \in [0, s^{(k)}(t)], \quad t \in [0, t_1],$$

by letting  $t \rightarrow t_1$ , the inequality (6.4) follows.

Next, we suppose that  $\lim_{t \rightarrow t_1^-} s^{(k)}(t)$  does not exist. We assume that

$$a \equiv \liminf_{t \rightarrow t_1^-} s^{(k)}(t) < \limsup_{t \rightarrow t_1^-} s^{(k)}(t) \equiv b.$$

Then  $\xi < a < b \leq 1$ . It is easy to see that  $(u^{(k)} - U)(x, t_1) = 0$  for all  $x \in [a, b]$  and  $(u^{(k)} - U)(x, t_1) > 0$  for all  $x \in [0, a)$ . If  $b = 1$ , then (6.4) follows immediately. Suppose that  $b < 1$ . For contradiction, we assume that  $(u^{(k)} - U)(x_0, t_1) < 0$  for some  $x_0 \in (b, 1)$ . Since  $\limsup_{t \rightarrow t_1^-} s^{(k)}(t) = b < x_0$ , there exists  $t_2 \in [0, t_1)$  such that  $s^{(k)}(t) < x_0$  for all  $t \in (t_2, t_1)$ . Hence

$$(u^{(k)} - U)(x, t) \leq 0, \quad \text{for all } (x, t) \in \{[x_0, 1] \times [t_2, t_1]\} \cup \{[s^{(k)}(t), 1] \times [0, t_2]\}.$$

It follows from the Hopf Boundary Point Lemma that  $(u^{(k)} - U)_x(1, t_1) > 0$ , a contradiction. Hence (6.4) follows.

Since  $u^{(k)}(\xi, t) > U(\xi)$  for all  $t \geq t_1$ , it follows from the maximum principle that  $u^{(k)}(x, t) \geq U(x)$  for all  $(x, t) \in [\xi, 1] \times [t_1, \infty)$ . In particular,  $u^{(n)}(1, t) \geq M$  for all  $t \geq T$  and  $n \geq k$ . Hence  $v(1, t) = \infty$  for all  $t \geq T$  and the lemma follows.  $\square$

Now, by the representation formula of solution of  $(B^{(n)})$ ,

$$\begin{aligned} u^{(n)}(x, t) &= \int_0^1 u^{(n)}(y, t_1) \Gamma(x, t; y, t_1) dy + \int_{t_1}^t f^{(n)}(u^{(n)}(1, \tau)) \Gamma(x, t; 1, \tau) d\tau \\ &\quad - \int_{t_1}^t u^{(n)}(1, \tau) \Gamma_y(x, t; 1, \tau) d\tau + \int_{t_1}^t u^{(n)}(0, \tau) \Gamma_y(x, t; 0, \tau) d\tau \\ &\quad + \int_{t_1}^t \int_0^1 g^{(n)}(u^{(n)}(y, \tau)) \Gamma(x, t; y, \tau) dy d\tau \end{aligned}$$

for  $x \in (0, 1)$  and  $t > t_1 \geq 0$ , where

$$\Gamma(x, t; y, \tau) = \frac{1}{\sqrt{4\pi(t-\tau)}} \exp\left\{-\frac{(x-y)^2}{4(t-\tau)}\right\},$$

and the jump relation of  $u^{(n)}(0, t)$

$$\begin{aligned} \frac{1}{2} u^{(n)}(0, t) &= \int_0^1 u^{(n)}(y, t_1) \Gamma(0, t; y, t_1) dy + \int_{t_1}^t f^{(n)}(u^{(n)}(1, \tau)) \Gamma(0, t; 1, \tau) d\tau \\ &\quad - \int_{t_1}^t u^{(n)}(1, \tau) \Gamma_y(0, t; 1, \tau) d\tau + \int_{t_1}^t \int_0^1 g^{(n)}(u^{(n)}(y, \tau)) \Gamma(0, t; y, \tau) dy d\tau \end{aligned}$$

for  $t > t_1 \geq 0$ , we conclude that  $v(x, t) \equiv \infty$  for all  $t > T$ . This proves that the blow-up is complete when  $q \geq 1$ .

#### REFERENCES

- [ 1 ] S. ANGENENT, The zeroset of a solution of a parabolic equation, *J. Reine Angew. Math.* 390 (1988), 79–96.
- [ 2 ] M. CHLEBÍK AND M. FILA, Some recent results on blow-up on the boundary for the heat equation, B. Bojarski et al. (ed.), *Proceedings of the Minisemester, Warsaw, Poland, (Warsaw, 1998)*, 61–71, Banach Center Publ. 52, Polish Acad. Sci., Warsaw, 2000.
- [ 3 ] A. DE PABLO, F. QUIRÓS AND J. D. ROSSI, Asymptotic simplification for a reaction-diffusion problem with a nonlinear boundary condition, *IMA J. Appl. Math.* 67 (2002), 69–98.
- [ 4 ] M. FILA AND J. FILO, Blow-up on the boundary: A survey S. Janeczko et al. (ed.), *Singularities and differential equations, (Warsaw, 1996)*, 67–78, Banach Center Publ. 33, Polish Acad. Sci., Warsaw, 1996.
- [ 5 ] M. FILA AND J.-S. GUO, Complete blow-up and incomplete quenching for the heat equation with a nonlinear boundary condition, *Nonlinear Anal.* 48 (2002), 995–1002.
- [ 6 ] M. FILA AND P. QUITNER, The blow-up rate for the heat equation with a nonlinear boundary condition, *Math. Methods Appl. Sci.* 14 (1991), 197–205.
- [ 7 ] A. FRIEDMAN AND B. MCLEOD, Blow-up of positive solutions of semilinear heat equations, *Indiana Univ. Math. J.* 34 (1985), 425–447.
- [ 8 ] Y. GIGA AND R. V. KOHN, Asymptotically self-similar blow-up of semilinear heat equations, *Comm. Pure Appl. Math.* 38 (1985), 297–319.
- [ 9 ] J.-S. GUO, On the quenching behavior of the solution of a semilinear parabolic equation, *J. Math. Anal. Appl.* 151 (1990), 58–79.
- [ 10 ] G. M. LIEBERMAN, *Second Order Parabolic Differential Equations*, World Scientific, Singapore, 1996.
- [ 11 ] Z. LIN AND M. WANG, The blow-up properties of solutions to semilinear heat equations with nonlinear boundary conditions, *Z. Angew. Math. Phys.* 50 (1999), 361–374.



- [12] M. WANG AND X. WANG, Existence of positive solutions to a nonlinear initial problem, *Nonlinear Anal.* 44 (2001), 1133–1136.
- [13] E. YANAGIDA, Uniqueness of positive radial solutions of  $\Delta u + g(r)u + h(r)u^p = 0$  in  $R^n$ , *Arch. Rational Mech. Anal.* 115 (1991), 257–274.
- [14] E. YANAGIDA, Structure of positive radial solutions of Matukuma's equation, *Japan J. Indust. Appl. Math.* 8 (1991), 165–173.

SHENG-CHEN FU:  
DEPARTMENT OF MATHEMATICAL SCIENCES  
NATIONAL CHENGCHI UNIVERSITY  
64 SEC. 2, ZHI-NAN ROAD  
TAIPEI 116  
TAIWAN

JONG-SHENQ GUO AND JE-CHIANG TSAI:  
DEPARTMENT OF MATHEMATICS  
NATIONAL TAIWAN NORMAL UNIVERSITY  
88, SEC. 4, TING CHOU ROAD  
TAIPEI 117  
TAIWAN